

Checklist of Yorkshire Coleoptera

Figures based on records on the YNU database in January 2011

Figures after family names are British species/Yorkshire species

<u>Taxon</u>	<u>VC61</u>	<u>VC62</u>	<u>VC63</u>	<u>VC64</u>	<u>VC65</u>	<u>VC Total</u>
<u>Gyrinidae [12/9]</u>						
Gyrinus aeratus Stephens, 1835	10	0	2	0	0	12
Gyrinus caspius Ménétrés, 1832	5	1	2	1	0	9
Gyrinus distinctus Aubé, 1838	1	0	2	0	0	3
Gyrinus marinus Gyllenhal, 1808	22	4	28	8	0	62
Gyrinus minutus Fabricius, 1798	0	2	3	0	0	5
Gyrinus paykulli Ochs, 1927	0	1	1	1	0	3
Gyrinus substriatus Stephens, 1828	22	14	79	20	4	139
Gyrinus urinator Illiger, 1807	0	3	1	0	0	4
Orectochilus villosus (Müller, O.F., 1776)	1	5	1	19	2	28
<u>Haliplidae [19/17]</u>						
Brychius elevatus (Panzer, 1793)	2	2	4	7	1	16
Peltodytes caesus (Duftschmid, 1805)	0	0	2	0	0	2
Haliplus (Haliplus) confinis Stephens, 1828	15	5	37	12	2	71
Haliplus (Haliplus) obliquus (Fabricius, 1787)	18	4	14	18	4	58
Haliplus (Neohalipus) lineatocollis (Marsham, 1802)	24	12	63	11	3	113
Haliplus (Haliplinus) apicalis Thomson, C.G., 1868	6	2	1	0	0	9
Haliplus (Haliplinus) fluviatilis Aubé, 1836	5	2	32	4	0	43
Haliplus (Haliplinus) heydeni Wehncke, 1875	2	2	6	3	0	13
Haliplus (Haliplinus) immaculatus Gerhardt, 1877	19	3	48	11	0	81
Haliplus (Haliplinus) lineolatus Mannerheim, 1844	13	3	25	9	0	50
Haliplus (Haliplinus) ruficollis (De Geer, 1774)	25	14	82	29	2	152
Haliplus (Haliplinus) sibiricus Motschulsky, 1860	44	22	34	20	4	124
Haliplus (Liaphlus) flavicollis Sturm, 1834	11	1	18	17	5	52
Haliplus (Liaphlus) fulvus (Fabricius, 1801)	7	3	7	7	1	25
Haliplus (Liaphlus) laminatus (Schaller, 1783)	0	0	9	6	0	15
Haliplus (Liaphlus) mucronatus Stephens, 1828	1	0	1	0	0	2
Haliplus (Liaphlus) variegatus Sturm, 1834	1	1	2	0	0	4
<u>Noteridae [2/2]</u>						
Noterus clavicornis (De Geer, 1774)	37	6	66	29	1	139
Noterus crassicornis (Müller, O.F., 1776)	5	2	10	2	0	19
<u>Paelobiidae [1/1]</u>						
Hygrobia hermanni (Fabricius, 1775)	17	4	19	6	1	47
<u>Dytiscidae [117/103]</u>						
Agabus (Acatodes) arcticus (Paykull, 1798)	0	0	15	8	4	27

Yorkshire Coleoptera checklist 2011

Agabus (Acatodes) congener (Thunberg, 1794)	4	2	7	10	4	27
Agabus (Acatodes) sturmii (Gyllenhal, 1808)	28	13	83	30	5	159
Agabus (Agabus) labiatus (Brahm, 1790)	6	2	2	1	0	11
Agabus (Agabus) uliginosus (Linnaeus, 1761)	9	9	7	12	1	38
Agabus (Agabus) undulatus (Schrank, 1776)	4	1	0	15	0	20
Agabus (Gaurodytes) affinis (Paykull, 1798)	4	2	6	3	0	15
Agabus (Gaurodytes) biguttatus (Olivier, 1795)	1	3	11	3	0	18
Agabus (Gaurodytes) bipustulatus (Linnaeus, 1767)	84	36	184	74	6	384
Agabus (Gaurodytes) conspersus (Marsham, 1802)	16	2	3	0	0	21
Agabus (Gaurodytes) didymus (Olivier, 1795)	21	8	35	13	3	80
Agabus (Gaurodytes) guttatus (Paykull, 1798)	0	12	64	21	3	100
Agabus (Gaurodytes) melanarius Aubé, 1837	1	3	1	0	0	5
Agabus (Gaurodytes) nebulosus (Forster, 1771)	33	16	65	32	2	148
Agabus (Gaurodytes) paludosus (Fabricius, 1801)	10	11	27	14	0	62
Agabus (Gaurodytes) striolatus (Gyllenhal, 1808)	0	0	1	0	0	1
Agabus (Gaurodytes) unguicularis (Thomson, C.G., 1867)	0	3	6	8	0	17
Ilybius aenescens Thomson, C.G., 1870	5	7	7	5	0	24
Ilybius ater (De Geer, 1774)	17	5	47	16	4	89
Ilybius chalconatus (Panzer, 1796)	10	17	12	10	1	50
Ilybius fenestratus (Fabricius, 1781)	7	2	18	4	0	31
Ilybius fuliginosus (Fabricius, 1792)	31	12	93	28	11	175
Ilybius guttiger (Gyllenhal, 1808)	8	2	7	7	0	24
Ilybius montanus (Stephens, 1828)	2	2	4	6	2	16
Ilybius quadriguttatus (Lacordaire, 1835)	16	2	24	8	0	50
Ilybius subaeneus Erichson, 1837	1	2	8	1	0	12
Platambus maculatus (Linnaeus, 1758)	6	12	30	21	6	75
Colymbetes fuscus (Linnaeus, 1758)	43	9	55	23	2	132
Rhantus (Nartus) grapii (Gyllenhal, 1808)	4	1	14	9	0	28
Rhantus (Rhantus) bistriatus (Bergsträsser, 1778)	0	0	1	0	0	1
Rhantus (Rhantus) exsoletus (Forster, 1771)	5	2	17	5	0	29
Rhantus (Rhantus) frontalis (Marsham, 1802)	1	0	0	1	0	2
Rhantus (Rhantus) suturalis (MacLeay, 1825)	2	1	8	6	0	17
Rhantus (Rhantus) suturellus (Harris, T.W., 1828)	13	6	9	5	0	33
Liopterus haemorrhoidalis (Fabricius, 1787)	14	0	19	11	0	44
Acilius canaliculatus (Nicolai, 1822)	3	0	26	0	0	29
Acilius sulcatus (Linnaeus, 1758)	9	12	27	9	0	57
Dytiscus circumcinctus Ahrens, 1811	3	1	12	9	0	25
Dytiscus circumflexus Fabricius, 1801	4	0	4	1	0	9
Dytiscus dimidiatus Bergsträsser, 1778	1	1	2	11	0	15
Dytiscus marginalis Linnaeus, 1758	22	11	73	14	3	123
Dytiscus semisulcatus Müller, O.F., 1776	6	4	13	6	1	30
Hydaticus continentalis Balfour-Browne, J., 1944	0	0	1	0	0	1
Hydaticus seminiger (De Geer, 1774)	0	0	0	4	0	4
Hydaticus transversalis (Pontoppidan, 1763)	0	0	2	7	0	9
Bidessus unistriatus (Goeze, 1777)	0	0	1	0	0	1
Hydroglyphus geminus (Fabricius, 1792)	4	1	12	4	0	21
Deronectes latus (Stephens, 1829)	0	4	2	2	0	8
Graptodytes bilineatus (Sturm, 1835)	6	0	0	0	0	6
Graptodytes granularis (Linnaeus, 1767)	3	0	7	5	0	15
Graptodytes pictus (Fabricius, 1787)	22	7	48	2	0	79

Yorkshire Coleoptera checklist 2011

Hydroporus angustatus Sturm, 1835	11	5	26	19	0	61
Hydroporus discretus Fairmaire & Brisout, 1859	10	6	16	11	1	44
Hydroporus erythrocephalus (Linnaeus, 1758)	25	3	22	27	0	77
Hydroporus ferrugineus Stephens, 1829	0	5	3	3	1	12
Hydroporus glabriusculus Aubé, 1838	0	0	0	1	0	1
Hydroporus gyllenhalii Schiödte, 1841	13	13	23	23	4	76
Hydroporus incognitus Sharp, 1869	1	5	13	5	0	24
Hydroporus longicornis Sharp, 1871	0	1	1	2	2	6
Hydroporus longulus Mulsant & Rey, 1861	0	2	5	2	3	12
Hydroporus marginatus (Duftschmid, 1805)	0	0	1	0	0	1
Hydroporus melanarius Sturm, 1835	5	11	12	12	3	43
Hydroporus memnonius Nicolai, 1822	14	17	27	18	3	79
Hydroporus morio Aubé, 1838	0	2	0	16	4	22
Hydroporus neglectus Schaum, 1845	3	1	6	3	0	13
Hydroporus nigrita (Fabricius, 1792)	7	11	20	16	5	59
Hydroporus obscurus Sturm, 1835	6	3	3	13	2	27
Hydroporus obsoletus Aubé, 1838	0	2	0	0	1	3
Hydroporus palustris (Linnaeus, 1761)	58	16	158	54	7	293
Hydroporus planus (Fabricius, 1781)	47	17	107	25	2	198
Hydroporus pubescens (Gyllenhal, 1808)	35	22	81	34	4	176
Hydroporus rufifrons (Müller, O.F., 1776)	0	4	3	8	0	15
Hydroporus scalesianus Stephens, 1828	0	0	3	3	0	6
Hydroporus striola (Gyllenhal, 1826)	6	3	14	13	0	36
Hydroporus tessellatus (Drapiez, 1819)	15	13	24	14	0	66
Hydroporus tristis (Paykull, 1798)	9	18	11	22	4	64
Hydroporus umbrosus (Gyllenhal, 1808)	9	10	13	13	1	46
Nebrioporus (Nebrioporus) assimilis (Paykull, 1798)	16	3	22	11	0	52
Nebrioporus (Nebrioporus) depressus (Fabricius, 1775)	0	0	0	1	0	1
Nebrioporus (Nebrioporus) elegans (Panzer, 1794)	7	2	21	16	3	49
Oreodytes davisii (Curtis, 1831)	0	6	6	9	7	28
Oreodytes sanmarkii (Sahlberg, C.R., 1826)	2	19	20	19	11	71
Oreodytes septentrionalis (Gyllenhal, 1826)	0	5	3	6	4	18
Porhydrus lineatus (Fabricius, 1775)	12	2	14	5	0	33
Scarodytes halensis (Fabricius, 1787)	5	0	5	5	0	15
Stictonectes lepidus (Olivier, 1795)	2	5	33	13	0	53
Stictotarsus duodecimpustulatus (Fabricius, 1792)	6	6	24	9	1	46
Stictotarsus multilineatus (Falkenström, 1922)	0	0	0	1	3	4
Suphrodytes dorsalis (Fabricius, 1787)	14	0	15	22	0	51
Hygrotus (Coelambus) confluens (Fabricius, 1787)	12	5	13	7	0	37
Hygrotus (Coelambus) impressopunctatus (Schaller, 1783)	27	6	64	25	1	123
Hygrotus (Coelambus) nigrolineatus (Steven, 1808)	0	0	1	1	0	2
Hygrotus (Coelambus) novemlineatus (Stephens, 1829)	0	2	0	0	3	5
Hygrotus (Coelambus) parallelogrammus (Ahrens, 1812)	8	0	0	0	0	8
Hygrotus (Hygrotus) decoratus (Gyllenhal, 1810)	2	2	0	7	0	11
Hygrotus (Hygrotus) inaequalis (Fabricius, 1777)	45	7	86	38	3	179
Hygrotus (Hygrotus) quinquelineatus (Zetterstedt, 1828)	0	0	3	1	0	4
Hygrotus (Hygrotus) versicolor (Schaller, 1783)	20	0	29	5	0	54
Hyphydrus ovatus (Linnaeus, 1761)	35	4	71	21	0	131
Laccornis oblongus (Stephens, 1835)	0	1	0	5	0	6
Laccophilus hyalinus (De Geer, 1774)	10	2	14	8	0	34

Yorkshire Coleoptera checklist 2011

<i>Laccophilus minutus</i> (Linnaeus, 1758)	26	11	56	36	1	130
<i>Laccophilus poecilus</i> Klug, 1834	0	0	4	0	0	4
<u>Carabidae [368/270]</u>						
<i>Cicindela campestris</i> Linnaeus, 1758	11	41	115	13	8	188
<i>Brachinus</i> (<i>Brachinus</i>) <i>crepitans</i> (Linnaeus, 1758)	0	0	1	0	0	1
<i>Carabus</i> (<i>Eucarabus</i>) <i>arvensis</i> Herbst, 1784	2	32	2	7	3	46
<i>Carabus</i> (<i>Carabus</i>) <i>granulatus</i> Linnaeus, 1758	20	3	26	4	0	53
<i>Carabus</i> (<i>Morphocarabus</i>) <i>monilis</i> Fabricius, 1792	4	8	9	12	0	33
<i>Carabus</i> (<i>Archicarabus</i>) <i>nemoralis</i> Müller, O.F., 1764	9	19	36	30	5	99
<i>Carabus</i> (<i>Autocarabus</i>) <i>auratus</i> Linnaeus, 1761	0	0	0	1	0	1
<i>Carabus</i> (<i>Autocarabus</i>) <i>cancellatus</i> Illiger, 1798	1	0	1	2	1	5
<i>Carabus</i> (<i>Hemicarabus</i>) <i>nitens</i> Linnaeus, 1758	2	40	12	9	18	81
<i>Carabus</i> (<i>Oreocarabus</i>) <i>glabratus</i> Paykull, 1790	0	10	1	0	3	14
<i>Carabus</i> (<i>Mesocarabus</i>) <i>problematicus</i> Herbst, 1786	6	33	70	19	18	146
<i>Carabus</i> (<i>Megodontus</i>) <i>violaceus</i> Linnaeus, 1758	8	16	39	19	15	97
<i>Cychrus caraboides</i> (Linnaeus, 1758)	2	15	82	31	11	141
<i>Leistus</i> (<i>Pogonophorus</i>) <i>rufomarginatus</i> (Duftschmid, 1812)	1	6	33	3	1	44
<i>Leistus</i> (<i>Pogonophorus</i>) <i>spinibarbis</i> (Fabricius, 1775)	7	5	23	4	2	41
<i>Leistus</i> (<i>Leistophorus</i>) <i>fulvibarbis</i> Dejean, 1826	15	7	75	35	5	137
<i>Leistus</i> (<i>Leistus</i>) <i>ferrugineus</i> (Linnaeus, 1758)	24	12	106	20	0	162
<i>Leistus</i> (<i>Leistus</i>) <i>terminatus</i> (Hellwig in Panzer, 1793)	16	27	72	44	7	166
<i>Nebria</i> (<i>Paranebria</i>) <i>livida</i> (Linnaeus, 1758)	12	6	0	0	0	18
<i>Nebria</i> (<i>Nebria</i>) <i>brevicollis</i> (Fabricius, 1792)	32	47	278	92	27	476
<i>Nebria</i> (<i>Nebria</i>) <i>salina</i> Fairmaire & Laboulbène, 1854	2	15	50	39	16	122
<i>Nebria</i> (<i>Boreonebria</i>) <i>rufescens</i> (Ström, 1768)	0	20	33	32	16	101
<i>Notiophilus aesthuans</i> Motschulsky, 1864	1	0	0	1	6	8
<i>Notiophilus aquaticus</i> (Linnaeus, 1758)	4	26	38	11	14	93
<i>Notiophilus biguttatus</i> (Fabricius, 1779)	76	57	298	102	34	567
<i>Notiophilus germinyi</i> Fauvel, 1863	3	23	12	20	12	70
<i>Notiophilus palustris</i> (Duftschmid, 1812)	17	12	50	12	2	93
<i>Notiophilus quadripunctatus</i> Dejean, 1826	4	3	4	0	0	11
<i>Notiophilus rufipes</i> Curtis, 1829	1	2	9	3	0	15
<i>Notiophilus substriatus</i> Waterhouse, G.R., 1833	11	3	30	15	1	60
<i>Blethisa multipunctata</i> (Linnaeus, 1758)	11	0	6	6	0	23
<i>Elaphrus</i> (<i>Elaphrus</i>) <i>cupreus</i> Duftschmid, 1812	24	11	86	36	3	160
<i>Elaphrus</i> (<i>Elaphrus</i>) <i>lapponicus</i> Gyllenhal, 1810	0	1	0	1	0	2
<i>Elaphrus</i> (<i>Elaphrus</i>) <i>uliginosus</i> Fabricius, 1792	1	0	0	0	0	1
<i>Elaphrus</i> (<i>Trichelaphrus</i>) <i>riparius</i> (Linnaeus, 1758)	41	13	38	36	5	133
<i>Loricera pilicornis</i> (Fabricius, 1775)	41	28	189	84	20	362
<i>Clivina collaris</i> (Herbst, 1784)	7	7	6	18	2	40
<i>Clivina fossor</i> (Linnaeus, 1758)	40	19	67	52	12	190
<i>Dyschirius</i> (<i>Dyschirius</i>) <i>angustatus</i> (Ahrens, 1830)	1	0	0	0	0	1
<i>Dyschirius</i> (<i>Dyschirius</i>) <i>thoracicus</i> (Rossi, 1790)	13	1	0	0	0	14
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>aeneus</i> (Dejean, 1825)	6	1	4	4	0	15
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>globosus</i> (Herbst, 1784)	27	24	66	10	6	133
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>luedersi</i> Wagner, 1915	16	3	18	1	0	38
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>nitidus</i> (Dejean, 1825)	3	1	0	0	0	4
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>politus</i> (Dejean, 1825)	7	1	3	1	1	13
<i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>salinus</i> Schaum, 1843	9	0	1	0	0	10

Yorkshire Coleoptera checklist 2011

<i>Brosicus cephalotes</i> (Linnaeus, 1758)	16	1	10	0	0	27
<i>Miscodera arctica</i> (Paykull, 1798)	2	13	8	9	7	39
<i>Aepus marinus</i> (Ström, 1783)	0	7	0	0	0	7
<i>Aepus robinii</i> (Laboulbène, 1849)	0	7	0	0	0	7
<i>Trechus</i> (<i>Epaphius</i>) <i>rivularis</i> (Gyllenhal, 1810)	0	0	0	8	0	8
<i>Trechus</i> (<i>Epaphius</i>) <i>secalis</i> (Paykull, 1790)	7	9	12	24	4	56
<i>Trechus</i> (<i>Trechus</i>) <i>fulvus</i> Dejean, 1831	0	1	0	0	0	1
<i>Trechus</i> (<i>Trechus</i>) <i>obtusus</i> Erichson, 1837	45	32	94	39	14	224
<i>Trechus</i> (<i>Trechus</i>) <i>quadristriatus</i> (Schrank, 1781)	58	13	149	29	13	262
<i>Trechus</i> (<i>Trechus</i>) <i>rubens</i> (Fabricius, 1792)	0	5	14	2	2	23
<i>Trechus</i> (<i>Trechus</i>) <i>subnotatus</i> Dejean, 1831	0	0	6	0	0	6
<i>Blemus discus</i> (Fabricius, 1792)	8	5	8	11	0	32
<i>Trechoblemus micros</i> (Herbst, 1784)	2	7	6	8	5	28
<i>Tachys</i> (<i>Paratachys</i>) <i>bistriatus</i> (Duftschmid, 1812)	1	0	0	0	0	1
<i>Tachys</i> (<i>Tachys</i>) <i>scutellaris</i> Stephens, 1828	0	1	0	0	0	1
<i>Elaphropus parvulus</i> (Dejean, 1831)	0	0	12	0	0	12
<i>Asaphidion curtum</i> (Heyden, 1870)	12	2	25	15	0	54
<i>Asaphidion flavipes</i> (Linnaeus, 1761)	2	1	1	5	0	9
<i>Asaphidion pallipes</i> (Duftschmid, 1812)	0	2	0	0	0	2
<i>Asaphidion stierlini</i> (Heyden, 1880)	10	0	13	5	0	28
<i>Ocys harpaloides</i> (Audinet-Serville, 1821)	23	17	83	40	7	170
<i>Ocys quinquestriatus</i> (Gyllenhal, 1810)	0	1	11	8	0	20
<i>Bracteon litorale</i> (Olivier, 1790)	2	13	1	10	4	30
<i>Bembidion</i> (<i>Neja</i>) <i>nigricorne</i> Gyllenhal, 1827	2	26	3	5	0	36
<i>Bembidion</i> (<i>Metallina</i>) <i>lampros</i> (Herbst, 1784)	65	49	247	83	12	456
<i>Bembidion</i> (<i>Metallina</i>) <i>properans</i> (Stephens, 1828)	12	25	51	14	2	104
<i>Bembidion</i> (<i>Princidium</i>) <i>punctulatum</i> Drapiez, 1821	0	15	3	23	7	48
<i>Bembidion</i> (<i>Testedium</i>) <i>bipunctatum</i> (Linnaeus, 1761)	1	1	1	2	1	6
<i>Bembidion</i> (<i>Eupetedromus</i>) <i>dentellum</i> (Thunberg, 1787)	32	6	69	32	5	144
<i>Bembidion</i> (<i>Notaphus</i>) <i>obliquum</i> Sturm, 1825	9	7	52	16	3	87
<i>Bembidion</i> (<i>Notaphus</i>) <i>varium</i> (Olivier, 1795)	47	1	23	1	0	72
<i>Bembidion</i> (<i>Plataphus</i>) <i>prasinum</i> (Duftschmid, 1812)	0	3	2	13	9	27
<i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>atrocaeruleum</i> (Stephens, 1828)	3	18	1	28	20	70
<i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>geniculatum</i> Heer, 1837/8	6	7	38	12	6	69
<i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>tibiale</i> (Duftschmid, 1812)	0	19	8	31	16	74
<i>Bembidion</i> (<i>Ocydromus</i>) <i>bruxellense</i> Wesmael, 1835	3	13	29	10	1	56
<i>Bembidion</i> (<i>Ocydromus</i>) <i>cruciatum</i> Dejean, 1831	13	8	2	0	2	25
<i>Bembidion</i> (<i>Ocydromus</i>) <i>decorum</i> (Zenker in Panzer, 1800)	0	17	2	21	10	50
<i>Bembidion</i> (<i>Ocydromus</i>) <i>deletum</i> Audinet-Serville, 1821	15	6	50	22	4	97
<i>Bembidion</i> (<i>Ocydromus</i>) <i>femoratum</i> Sturm, 1825	5	5	16	10	2	38
<i>Bembidion</i> (<i>Ocydromus</i>) <i>fluviatile</i> Dejean, 1831	0	0	0	3	0	3
<i>Bembidion</i> (<i>Ocydromus</i>) <i>lunatum</i> (Duftschmid, 1812)	11	4	11	6	0	32
<i>Bembidion</i> (<i>Ocydromus</i>) <i>maritimum</i> (Stephens, 1835)	7	2	4	0	0	13
<i>Bembidion</i> (<i>Ocydromus</i>) <i>monticola</i> Sturm, 1825	0	4	2	9	3	18
<i>Bembidion</i> (<i>Ocydromus</i>) <i>saxatile</i> Gyllenhal, 1827	2	9	2	1	1	15
<i>Bembidion</i> (<i>Ocydromus</i>) <i>stephensii</i> Crotch, 1866	10	8	7	5	0	30
<i>Bembidion</i> (<i>Ocydromus</i>) <i>testaceum</i> (Duftschmid, 1812)	0	1	0	2	2	5
<i>Bembidion</i> (<i>Ocydromus</i>) <i>tetracolum</i> Say, 1825	31	26	103	61	30	251
<i>Bembidion</i> (<i>Nepha</i>) <i>illigeri</i> Netolitzky, 1914	33	9	29	10	2	83
<i>Bembidion</i> (<i>Sinechostictus</i>) <i>stomoides</i> Dejean, 1831	1	2	0	14	4	21

Yorkshire Coleoptera checklist 2011

Bembidion (Semicampa) gilvipes Sturm, 1825	46	19	19	16	0	100
Bembidion (Semicampa) schuppelii Dejean, 1831	0	0	1	1	0	2
Bembidion (Diplocampa) assimile Gyllenhal, 1810	34	2	60	1	1	98
Bembidion (Diplocampa) clarkii (Dawson, 1849)	18	11	37	20	0	86
Bembidion (Diplocampa) fumigatum (Duftschmid, 1812)	24	1	38	2	0	65
Bembidion (Emphanes) minimum (Fabricius, 1792)	43	0	0	1	0	44
Bembidion (Emphanes) normannum Dejean, 1831	6	0	0	0	0	6
Bembidion (Bembidion) humerale Sturm, 1825	0	0	26	0	0	26
Bembidion (Bembidion) quadrimaculatum (Linnaeus, 1761)	29	9	55	32	2	127
Bembidion (Bembidion) quadripustulatum Audinet-Serville, 1821	0	0	1	0	0	1
Bembidion (Trepanedoris) doris (Panzer, 1796)	17	3	16	18	0	54
Bembidion (Trepanes) articulatum (Panzer, 1795)	7	0	8	1	0	16
Bembidion (Phyla) obtusum Audinet-Serville, 1821	47	15	64	30	1	157
Bembidion (Philochthus) aeneum Germar, 1824	52	24	94	45	5	220
Bembidion (Philochthus) biguttatum (Fabricius, 1779)	64	24	83	42	2	215
Bembidion (Philochthus) guttula (Fabricius, 1792)	60	32	194	75	8	369
Bembidion (Philochthus) iricolor Bedel, 1879	20	0	4	0	0	24
Bembidion (Philochthus) lunulatum (Geoffroy in Fourcroy, 1785)	13	2	54	25	0	94
Bembidion (Philochthus) mannerheimii Sahlberg, C.R., 1827	8	20	20	19	5	72
Pogonus chalceus (Marsham, 1802)	10	0	0	0	0	10
Pogonus littoralis (Duftschmid, 1812)	1	0	0	0	0	1
Patrobus assimilis Chaudoir, 1844	1	14	2	16	12	45
Patrobus atrorufus (Ström, 1768)	6	11	29	41	10	97
Patrobus septentrionis Dejean, 1828	2	0	2	0	0	4
Stomis pumicatus (Panzer, 1795)	16	12	32	18	1	79
Poecilus cupreus (Linnaeus, 1758)	11	11	47	5	1	75
Poecilus lepidus (Leske, 1785)	3	18	0	0	0	21
Poecilus versicolor (Sturm, 1824)	14	20	46	10	1	91
Pterostichus (Pterostichus) cristatus (Dufour, 1820)	0	0	2	6	7	15
Pterostichus (Steropus) aethiops (Panzer, 1796)	0	3	0	4	3	10
Pterostichus (Steropus) madidus (Fabricius, 1775)	29	48	378	124	39	618
Pterostichus (Pediis) longicollis (Duftschmid, 1812)	4	0	0	2	0	6
Pterostichus (Adelosia) macer (Marsham, 1802)	12	9	6	4	1	32
Pterostichus (Platysma) niger (Schaller, 1783)	40	21	216	55	16	348
Pterostichus (Bothriopterus) adstrictus Eschscholtz, 1823	3	15	13	18	11	60
Pterostichus (Bothriopterus) oblongopunctatus (Fabricius, 1787)	9	23	21	4	0	57
Pterostichus (Bothriopterus) quadrifoveolatus Letzner, 1852	7	2	36	11	0	56
Pterostichus (Omaseus) melanarius (Illiger, 1798)	37	15	97	62	10	221
Pterostichus (Pseudomaseus) anthracinus (Panzer, 1795)	15	0	5	0	0	20
Pterostichus (Pseudomaseus) gracilis (Dejean, 1828)	6	0	2	6	0	14
Pterostichus (Pseudomaseus) minor (Gyllenhal, 1827)	24	17	91	27	0	159
Pterostichus (Pseudomaseus) nigrita (Paykull, 1790)	25	34	152	50	5	266
Pterostichus (Pseudomaseus) rhaeticus Heer, 1837/8	2	7	23	12	3	47
Pterostichus (Lagarus) vernalis (Panzer, 1795)	13	9	32	25	0	79
Pterostichus (Argutor) diligens (Sturm, 1824)	33	44	242	84	12	415
Pterostichus (Argutor) strenuus (Panzer, 1796)	92	40	329	119	19	599
Abax parallelepipedus (Piller & Mitterpacher, 1783)	10	11	130	53	9	213
Platyderus depressus (Audinet-Serville, 1821)	8	1	3	0	0	12
Synuchus vivalis (Illiger, 1798)	6	6	24	11	4	51
Calathus (Amphigynus) rotundicollis Dejean, 1828	6	2	37	11	0	56

Yorkshire Coleoptera checklist 2011

Calathus (Calathus) ambiguus (Paykull, 1790)	4	3	6	0	0	13
Calathus (Calathus) erratus (Sahlberg, C.R., 1827)	4	9	7	2	0	22
Calathus (Calathus) fuscipes (Goeze, 1777)	20	9	62	22	8	121
Calathus (Calathus) melanocephalus (Linnaeus, 1758)	18	1	61	10	7	97
Calathus (Calathus) micropterus (Duftschmid, 1812)	0	13	5	14	14	46
Calathus (Calathus) mollis (Marsham, 1802)	24	6	1	0	0	31
Sphodrus leucophthalmus (Linnaeus, 1758)	0	2	3	0	0	5
Laemostenus (Laemostenus) complanatus (Dejean, 1828)	1	1	1	1	0	4
Laemostenus (Pristonychus) terricola (Herbst, 1784)	7	8	8	4	0	27
Olisthopus rotundatus (Paykull, 1790)	4	20	52	14	5	95
Oxypselaphus obscurus (Herbst, 1784)	23	7	152	47	4	233
Paranchus albipes (Fabricius, 1796)	33	40	180	74	29	356
Batenus livens (Gyllenhal, 1810)	6	0	14	0	0	20
Platynus assimilis (Paykull, 1790)	6	17	148	41	8	220
Anchomenus dorsalis (Pontoppidan, 1763)	42	10	81	27	1	161
Sericoda quadripunctata (De Geer, 1774)	0	1	0	0	0	1
Agonum (Europhilus) fuliginosum (Panzer, 1809)	66	21	226	73	20	406
Agonum (Europhilus) gracile Sturm, 1824	14	9	79	22	3	127
Agonum (Europhilus) micans Nicolai, 1822	7	2	21	11	1	42
Agonum (Europhilus) piceum (Linnaeus, 1758)	12	5	20	5	0	42
Agonum (Europhilus) thoreyi Dejean, 1828	37	5	87	21	2	152
Agonum (Agonum) emarginatum (Gyllenhal, 1827)	25	7	24	12	3	71
Agonum (Agonum) ericeti (Panzer, 1809)	0	7	11	6	2	26
Agonum (Agonum) gracilipes (Duftschmid, 1812)	2	0	0	0	0	2
Agonum (Agonum) marginatum (Linnaeus, 1758)	27	8	89	16	4	144
Agonum (Agonum) muelleri (Herbst, 1784)	8	14	46	34	5	107
Agonum (Agonum) nigrum Dejean, 1828	0	0	1	0	0	1
Agonum (Agonum) versutum Sturm, 1824	10	0	2	0	0	12
Agonum (Agonum) viduum (Panzer, 1796)	6	0	7	7	0	20
Amara (Zezea) plebeja (Gyllenhal, 1810)	31	26	165	49	5	276
Amara (Amara) aenea (De Geer, 1774)	21	34	178	23	7	263
Amara (Amara) anthobia Villa & Villa, 1833	0	0	2	0	0	2
Amara (Amara) communis (Panzer, 1797)	4	8	41	17	2	72
Amara (Amara) convexior Stephens, 1828	0	3	4	1	0	8
Amara (Amara) curta Dejean, 1828	0	3	1	0	0	4
Amara (Amara) eurynota (Panzer, 1796)	6	5	23	3	1	38
Amara (Amara) famelica Zimmermann, 1832	1	3	0	0	0	4
Amara (Amara) familiaris (Duftschmid, 1812)	38	24	136	40	6	244
Amara (Amara) lucida (Duftschmid, 1812)	7	1	2	1	0	11
Amara (Amara) lunicollis Schiödte, 1837	4	12	75	14	1	106
Amara (Amara) ovata (Fabricius, 1792)	9	4	17	12	2	44
Amara (Amara) similata (Gyllenhal, 1810)	22	11	75	21	3	132
Amara (Amara) sprete Dejean, 1831	0	2	0	0	0	2
Amara (Amara) tibialis (Paykull, 1798)	15	2	25	1	0	43
Amara (Celia) bifrons (Gyllenhal, 1810)	8	0	12	8	1	29
Amara (Celia) fusca Dejean, 1828	0	0	1	0	0	1
Amara (Celia) praetermissa (Sahlberg, C.R., 1827)	2	2	9	0	1	14
Amara (Bradytus) apricaria (Paykull, 1790)	8	5	45	9	9	76
Amara (Bradytus) consularis (Duftschmid, 1812)	4	2	2	2	0	10
Amara (Bradytus) fulva (Müller, O.F., 1776)	11	0	9	0	2	22

Yorkshire Coleoptera checklist 2011

<i>Amara (Percosia) equestris</i> (Duftschmid, 1812)	0	1	0	0	0	1
<i>Curtonotus aulicus</i> (Panzer, 1796)	24	17	74	30	3	148
<i>Curtonotus convexiusculus</i> (Marshall, 1802)	26	3	14	7	0	50
<i>Harpalus (Pseudoophonus) calceatus</i> (Duftschmid, 1812)	1	0	0	0	0	1
<i>Harpalus (Pseudoophonus) rufipes</i> (De Geer, 1774)	38	13	83	28	4	166
<i>Harpalus (Harpalus) affinis</i> (Schrank, 1781)	30	12	76	21	4	143
<i>Harpalus (Harpalus) anxius</i> (Duftschmid, 1812)	1	0	0	0	0	1
<i>Harpalus (Harpalus) attenuatus</i> Stephens, 1828	1	0	0	0	0	1
<i>Harpalus (Harpalus) froelichii</i> Sturm, 1818	2	0	0	0	0	2
<i>Harpalus (Harpalus) latus</i> (Linnaeus, 1758)	9	11	10	5	2	37
<i>Harpalus (Harpalus) neglectus</i> Audinet-Serville, 1821	1	0	0	0	0	1
<i>Harpalus (Harpalus) rubripes</i> (Duftschmid, 1812)	3	2	12	4	0	21
<i>Harpalus (Harpalus) rufipalpis</i> Sturm, 1818	0	0	9	1	0	10
<i>Harpalus (Harpalus) servus</i> (Duftschmid, 1812)	1	0	1	0	0	2
<i>Harpalus (Harpalus) smaragdinus</i> (Duftschmid, 1812)	0	1	0	0	0	1
<i>Harpalus (Harpalus) tardus</i> (Panzer, 1796)	26	1	6	1	0	34
<i>Ophonus (Ophonus) ardosiacus</i> Lutshnik, 1922	5	1	0	0	0	6
<i>Ophonus (Metophonus) puncticeps</i> Stephens, 1828	3	0	2	0	0	5
<i>Ophonus (Metophonus) puncticollis</i> (Paykull, 1798)	2	0	2	0	0	4
<i>Ophonus (Metophonus) rufibarbis</i> (Fabricius, 1792)	32	9	41	8	5	95
<i>Ophonus (Metophonus) rupicola</i> (Sturm, 1818)	1	0	0	0	0	1
<i>Ophonus (Metophonus) schaubergerianus</i> (Puel, 1937)	3	3	1	1	0	8
<i>Anisodactylus binotatus</i> (Fabricius, 1787)	2	2	53	4	0	61
<i>Dicheirotichus gustavii</i> Crotch, 1871	15	0	0	0	0	15
<i>Dicheirotichus obsoletus</i> (Dejean, 1829)	1	0	0	0	0	1
<i>Trichocellus cognatus</i> (Gyllenhal, 1827)	6	22	29	15	6	78
<i>Trichocellus placidus</i> (Gyllenhal, 1827)	37	23	130	60	4	254
<i>Bradycellus caucasicus</i> (Chaudoir, 1846)	2	6	12	10	7	37
<i>Bradycellus distinctus</i> (Dejean, 1829)	0	2	0	0	0	2
<i>Bradycellus harpalinus</i> (Audinet-Serville, 1821)	13	26	133	36	4	212
<i>Bradycellus ruficollis</i> (Stephens, 1828)	4	34	52	14	8	112
<i>Bradycellus sharpi</i> Joy, 1912	2	5	12	5	0	24
<i>Bradycellus verbasci</i> (Duftschmid, 1812)	8	4	56	11	4	83
<i>Stenolophus mixtus</i> (Herbst, 1784)	25	6	96	25	1	153
<i>Acupalpus brunnipes</i> (Sturm, 1825)	0	0	1	0	0	1
<i>Acupalpus dubius</i> Schilsky, 1888	19	23	173	32	1	248
<i>Acupalpus exiguus</i> Dejean, 1829	1	0	5	0	0	6
<i>Acupalpus meridianus</i> (Linnaeus, 1761)	5	0	12	2	0	19
<i>Acupalpus parvulus</i> (Sturm, 1825)	21	3	43	8	0	75
<i>Anthracus consputus</i> (Duftschmid, 1812)	7	1	4	4	0	16
<i>Chlaenius nigricornis</i> (Fabricius, 1787)	14	2	1	3	1	21
<i>Chlaenius tristis</i> (Schaller, 1783)	1	0	0	0	0	1
<i>Chlaenius vestitus</i> (Paykull, 1790)	19	1	2	0	0	22
<i>Oodes helopioides</i> (Fabricius, 1792)	1	1	0	1	0	3
<i>Licinus depressus</i> (Paykull, 1790)	9	1	1	0	0	11
<i>Licinus punctatulus</i> (Fabricius, 1792)	0	1	0	0	0	1
<i>Badister (Badister) bullatus</i> (Schrank, 1798)	43	10	85	25	1	164
<i>Badister (Badister) unipustulatus</i> Bonelli, 1813	4	0	11	0	0	15
<i>Badister (Trimorphus) sodalis</i> (Duftschmid, 1812)	1	3	17	4	0	25
<i>Panagaeus bipustulatus</i> (Fabricius, 1775)	13	0	0	0	0	13

Yorkshire Coleoptera checklist 2011

Panagaeus cruxmajor (Linnaeus, 1758)	1	0	2	0	0	3
Perigona nigriceps (Dejean, 1831)	1	0	0	2	0	3
Lebia (Lamprias) chlorocephala (Hoffmann, J., 1803)	3	7	2	0	0	12
Lebia (Lamprias) cyanocephala (Linnaeus, 1758)	0	0	0	1	0	1
Demetrias (Demetrias) atricapillus (Linnaeus, 1758)	77	19	113	23	1	233
Demetrias (Demetrias) monostigma Samouelle, 1819	7	1	0	0	0	8
Cymindis (Tarulus) vaporariorum (Linnaeus, 1758)	0	9	3	0	0	12
Paradromius linearis (Olivier, 1795)	104	35	207	50	7	403
Paradromius longiceps Dejean, 1826	8	0	7	0	0	15
Dromius agilis (Fabricius, 1787)	2	3	11	10	0	26
Dromius angustus Brullé, 1834	1	7	2	2	0	12
Dromius meridionalis Dejean, 1825	11	4	7	7	0	29
Dromius quadrimaculatus (Linnaeus, 1758)	9	20	47	23	1	100
Calodromius spilotus (Illiger, 1798)	17	13	32	9	1	72
Philorhizus melanocephalus (Dejean, 1825)	65	11	125	26	0	227
Philorhizus notatus (Stephens, 1827)	12	4	0	1	0	17
Philorhizus quadrisignatus (Dejean, 1825)	0	0	1	0	0	1
Philorhizus sigma (Rossi, 1790)	0	1	6	7	0	14
Microlestes maurus (Sturm, 1827)	18	0	1	0	0	19
Lionychus quadrillum (Duftschmid, 1812)	0	1	0	0	0	1
Syntomus foveatus (Geoffroy in Fourcroy, 1785)	21	8	15	1	0	45
Syntomus obscuroguttatus (Duftschmid, 1812)	0	0	1	0	0	1
Syntomus truncatellus (Linnaeus, 1761)	8	1	5	1	0	15
Odacantha melanura (Linnaeus, 1767)	0	0	1	0	0	1

Helophoridae [20/19]

Helophorus (Empleurus) nubilus Fabricius, 1777	4	3	7	9	0	23
Helophorus (Empleurus) porculus Bedel, 1881	0	4	0	2	0	6
Helophorus (Empleurus) rufipes (Bosc d'Antic, 1791)	1	0	0	0	0	1
Helophorus (Cyphelophorus) tuberculatus Gyllenhal, 1808	0	10	2	2	0	14
Helophorus (Trichohelophorus) alternans Gené 1836	0	0	0	1	0	1
Helophorus (Megahelophorus) aequalis Thomson, C.G., 1868	25	8	74	24	4	135
Helophorus (Megahelophorus) grandis Illiger, 1798	22	24	98	18	7	169
Helophorus (Atracthelophorus) arvernicus Mulsant, 1846	1	7	3	2	7	20
Helophorus (Atracthelophorus) brevipalpis Bedel, 1881	44	44	187	54	26	355
Helophorus (Helophorus) dorsalis (Marsham, 1802)	4	1	1	0	0	6
Helophorus (Helophorus) flavipes Fabricius, 1792	7	19	37	19	7	89
Helophorus (Helophorus) fulgidicollis Motschulsky, 1860	6	0	0	0	0	6
Helophorus (Helophorus) granularis (Linnaeus, 1760)	3	1	4	3	0	11
Helophorus (Helophorus) griseus Herbst, 1793	9	1	7	1	0	18
Helophorus (Helophorus) longitarsis Wollaston, 1864	2	0	0	0	0	2
Helophorus (Helophorus) minutus Fabricius, 1775	41	8	82	31	5	167
Helophorus (Helophorus) nanus Sturm, 1836	11	0	2	3	0	16
Helophorus (Helophorus) obscurus Mulsant, 1844	11	5	24	7	1	48
Helophorus (Helophorus) strigifrons Thomson, C.G., 1868	1	0	1	3	0	5

Georissidae [1/1]

Georissus crenulatus (Rossi, 1794)	12	6	0	1	0	19
------------------------------------	----	---	---	---	---	----

Hydrochidae [7/3]

Hydrochus angustatus Germar, 1824	1	0	0	1	0	2
Hydrochus brevis (Herbst, 1793)	1	0	1	0	0	2
Hydrochus elongatus (Schaller, 1783)	1	1	7	3	0	12

Spercheidae [1/1]

Spercheus emarginatus (Schaller, 1783)	0	1	0	2	0	3
--	---	---	---	---	---	---

Hydrophilidae [70/62]

Anacaena bipustulata (Marsham, 1802)	1	0	3	0	0	4
Anacaena globulus (Paykull, 1798)	64	41	289	66	27	487
Anacaena limbata (Fabricius, 1792)	53	6	89	34	1	183
Anacaena lutescens (Stephens, 1829)	12	4	53	19	2	90
Berosus (Berosus) affinis Brullé, 1835	3	1	3	1	0	8
Berosus (Berosus) luridus (Linnaeus, 1761)	3	0	5	0	0	8
Berosus (Berosus) signaticollis (Charpentier, 1825)	1	0	0	2	0	3
Berosus (Enoplurus) fulvus Kuwert, 1888	0	1	0	0	0	1
Chaetarthria seminulum (Herbst, 1797)	10	11	13	8	0	42
Chaetarthria simillima Vorst & Cuppen, 2003	3	0	0	0	0	3
Cymbiodyta marginellus (Fabricius, 1792)	24	3	19	15	0	61
Enochrus Thomson, C.G., 1859	0	0	1	0	0	1
Enochrus affinis (Thunberg, 1794)	7	5	5	4	0	21
Enochrus bicolor (Fabricius, 1792)	8	1	3	0	0	12
Enochrus coarctatus (Gredler, 1863)	2	3	5	6	0	16
Enochrus fuscipennis (Thomson, C.G., 1884)	2	0	1	0	0	3
Enochrus halophilus (Bedel, 1878)	6	0	0	0	0	6
Enochrus melanocephalus (Olivier, 1792)	3	3	11	5	0	22
Enochrus ochropterus (Marsham, 1802)	3	1	6	3	0	13
Enochrus quadripunctatus (Herbst, 1797)	1	1	0	1	0	3
Enochrus testaceus (Fabricius, 1801)	13	3	16	10	0	42
Helochares lividus (Forster, 1771)	17	0	12	7	0	36
Helochares obscurus (Müller, O.F., 1776)	3	1	5	0	0	9
Helochares punctatus Sharp, 1869	6	4	25	5	0	40
Hydrobius fuscipes (Linnaeus, 1758)	68	27	146	32	14	287
Hydrochara caraboides (Linnaeus, 1758)	0	0	2	2	0	4
Hydrophilus piceus (Linnaeus, 1758)	0	0	3	0	0	3
Laccobius atratus Rottenberg, 1874	1	2	0	0	1	4
Laccobius bipunctatus (Fabricius, 1775)	11	11	51	17	3	93
Laccobius colon (Stephens, 1829)	10	2	29	15	0	56
Laccobius minutus (Linnaeus, 1758)	14	6	26	7	4	57
Laccobius sinuatus Motschulsky, 1849	9	1	11	3	0	24
Laccobius striatulus (Fabricius, 1801)	0	5	5	6	3	19
Laccobius ytenensis Sharp, 1910	0	0	1	0	0	1
Coelostoma orbiculare (Fabricius, 1775)	5	12	11	9	1	38
Cercyon (Cercyon) bifenestratus Küster, 1851	0	0	3	0	0	3
Cercyon (Cercyon) convexiusculus Stephens, 1829	25	2	44	21	3	95
Cercyon (Cercyon) depressus Stephens, 1829	3	0	1	0	0	4
Cercyon (Cercyon) granarius Erichson, 1837	2	1	0	0	0	3
Cercyon (Cercyon) haemorrhoidalis (Fabricius, 1775)	35	13	92	22	13	175
Cercyon (Cercyon) impressus (Sturm, 1807)	14	13	85	20	13	145

Yorkshire Coleoptera checklist 2011

Cercyon (Cercyon) lateralis (Marsham, 1802)	27	25	93	18	13	176
Cercyon (Cercyon) littoralis (Gyllenhal, 1808)	16	6	0	0	0	22
Cercyon (Cercyon) marinus Thomson, C.G., 1853	18	1	17	2	1	39
Cercyon (Cercyon) melanocephalus (Linnaeus, 1758)	35	18	152	34	14	253
Cercyon (Cercyon) nigriceps (Marsham, 1802)	3	4	4	2	0	13
Cercyon (Cercyon) obsoletus (Gyllenhal, 1808)	4	2	16	2	0	24
Cercyon (Cercyon) pygmaeus (Illiger, 1801)	11	8	22	8	3	52
Cercyon (Cercyon) quisquilius (Linnaeus, 1761)	7	5	23	3	12	50
Cercyon (Cercyon) terminatus (Marsham, 1802)	11	3	10	2	1	27
Cercyon (Cercyon) tristis (Illiger, 1801)	15	7	8	17	1	48
Cercyon (Cercyon) unipunctatus (Linnaeus, 1758)	15	10	32	6	6	69
Cercyon (Paracercyon) analis (Paykull, 1798)	26	5	65	8	0	104
Cercyon (Dicyrtocercyon) ustulatus (Preyssl, 1790)	11	5	26	6	2	50
Megasternum concinnum (Marsham, 1802)	86	31	239	76	32	464
Cryptopleurum crenatum (Kugelann, 1794)	0	0	0	3	0	3
Cryptopleurum minutum (Fabricius, 1775)	29	14	100	18	8	169
Cryptopleurum subtile Sharp, 1884	3	1	1	1	0	6
Sphaeridium bipustulatum Fabricius, 1781	8	5	14	10	1	38
Sphaeridium lunatum Fabricius, 1792	15	12	34	4	1	66
Sphaeridium marginatum Fabricius, 1787	1	0	2	0	0	3
Sphaeridium scarabaeoides (Linnaeus, 1758)	9	3	38	8	6	64

Sphaeritidae [1/1]

Sphaerites glabratus (Fabricius, 1792)	0	4	0	1	0	5
--	---	---	---	---	---	---

Histeridae [51/32]

Abraeus granulum Erichson, 1839	0	0	2	0	0	2
Abraeus perpusillus (Marsham, 1802)	3	7	5	1	1	17
Plegaderus vulneratus (Panzer, 1797)	0	0	2	1	0	3
Acritus nigricornis (Hoffmann, J., 1803)	4	3	6	1	0	14
Aeletes atomarius (Aubé, 1842)	0	1	1	0	0	2
Saprinus aeneus (Fabricius, 1775)	3	2	5	0	0	10
Saprinus planiusculus Motschulsky, 1849	1	0	0	0	0	1
Saprinus semistriatus (Scriba, 1790)	9	9	15	5	1	39
Saprinus virescens (Paykull, 1798)	1	0	0	2	0	3
Hypocaccus (Hypocaccus) rugiceps (Duftschmid, 1805)	1	0	0	0	0	1
Hypocaccus (Hypocaccus) rugifrons (Paykull, 1798)	0	1	1	0	0	2
Hypocaccus (Baeckmanniolus) dimidiatus (Illiger, 1807)	1	2	0	0	0	3
Gnathoncus nannetensis (Marseul, 1862)	3	5	1	2	0	11
Gnathoncus rotundatus (Kugelann, 1792)	1	0	1	2	0	4
Myrmetes paykulli Kanaar, 1979	0	5	0	0	0	5
Dendrophilus punctatus (Herbst, 1792)	0	1	0	0	0	1
Dendrophilus pygmaeus (Linnaeus, 1758)	0	0	2	0	0	2
Kissister minimus (Aubé, 1850)	0	0	1	0	0	1
Carcinops pumilio (Erichson, 1834)	8	2	14	3	1	28
Paromalus flavicornis (Herbst, 1792)	0	4	9	3	0	16
Paromalus parallelepipedus (Herbst, 1792)	0	0	1	0	0	1
Onthophilus striatus (Forster, 1771)	18	10	60	10	1	99
Margarinotus (Ptomister) brunneus (Fabricius, 1775)	14	6	22	6	1	49
Margarinotus (Ptomister) merdarius (Hoffmann, J., 1803)	2	0	14	0	0	16

Yorkshire Coleoptera checklist 2011

Margarinotus (Ptomister) striola (Sahlberg, C.R., 1819)	2	7	15	10	5	39
Margarinotus (Paralister) neglectus (Germar, 1813)	0	1	0	1	0	2
Margarinotus (Paralister) purpurascens (Herbst, 1792)	7	1	2	0	0	10
Margarinotus (Paralister) ventralis (Marseul, 1854)	8	6	24	5	2	45
Margarinotus (Promethister) marginatus (Erichson, 1834)	4	2	1	0	0	7
Hister unicolor Linnaeus, 1758	5	4	7	0	2	18
Atholus bimaculatus (Linnaeus, 1758)	2	1	10	2	0	15
Atholus duodecimstriatus (Schrank, 1781)	3	4	16	3	1	27

Hydraenidae [30/20]

Hydraena britteni Joy, 1907	3	1	16	7	1	28
Hydraena flavipes Sturm, 1836	0	2	2	0	0	4
Hydraena gracilis Germar, 1824	0	8	4	9	8	29
Hydraena nigrita Germar, 1824	1	5	7	9	4	26
Hydraena palustris Erichson, 1837	1	1	2	2	1	7
Hydraena pulchella Germar, 1824	0	1	2	0	0	3
Hydraena pygmaea Waterhouse, G.R., 1833	0	3	2	2	0	7
Hydraena riparia Kugelann, 1794	8	5	11	10	1	35
Hydraena rufipes Curtis, 1830	0	3	0	3	2	8
Hydraena testacea Curtis, 1830	4	3	10	1	0	18
Limnebius aluta Bedel, 1881	1	0	1	3	0	5
Limnebius nitidus (Marsham, 1802)	0	2	7	0	0	9
Limnebius papposus Mulsant, 1844	1	0	5	1	0	7
Limnebius truncatellus (Thunberg, 1794)	4	14	34	6	5	63
Enicocerus exsculptus (Germar, 1824)	3	5	2	7	8	25
Ochthebius (Asiobates) auriculatus Rey, 1885	3	0	0	0	0	3
Ochthebius (Asiobates) bicolon Germar, 1824	3	4	7	8	1	23
Ochthebius (Asiobates) dilatatus Stephens, 1829	20	3	8	2	0	33
Ochthebius (Homalochthebius) minimus (Fabricius, 1792)	49	15	37	24	5	130
Ochthebius (Ochthebius) marinus (Paykull, 1798)	10	1	0	0	0	11

Ptiliidae [75/57]

Nossidium pilosellum (Marsham, 1802)	2	0	0	0	0	2
Ptenidium (Gressnerium) gressneri Erichson, 1845	0	0	0	1	0	1
Ptenidium (Matthewsium) laevigatum Erichson, 1845	8	8	19	8	1	44
Ptenidium (Matthewsium) turgidum Thomson, C.G., 1855	0	2	0	1	0	3
Ptenidium (Wankowizium) intermedium Wankowicz, 1869	5	1	9	5	1	21
Ptenidium (Ptenidium) formicetorum Kraatz, 1851	0	0	1	0	0	1
Ptenidium (Ptenidium) fuscicorne Erichson, 1845	13	7	27	4	1	52
Ptenidium (Ptenidium) punctatum (Gyllenhal, 1827)	4	5	7	0	2	18
Ptenidium (Ptenidium) pusillum (Gyllenhal, 1808)	27	5	75	22	8	137
Ptenidium (Gillmeisterium) nitidum (Heer, 1841)	20	22	93	24	15	174
Euryptilium gillmeisteri Flach, 1889	0	0	0	1	0	1
Euryptilium saxonicum (Gillmeister, 1845)	0	0	2	0	0	2
Ptiliola kunzei (Heer, 1841)	1	3	10	3	0	17
Ptiliolium (Ptiliolium) fuscum (Erichson, 1845)	4	14	13	8	4	43
Ptiliolium (Ptiliolium) marginatum (Aubé, 1850)	0	0	1	0	0	1
Ptiliolium (Ptiliolium) sahlbergi (Flach, 1888)	0	0	0	1	0	1
Ptiliolium (Ptiliolium) spencei (Allibert, 1844)	0	4	13	4	2	23
Ptiliolium (Euptilium) schwarzi (Flach, 1887)	0	0	2	0	0	2

Yorkshire Coleoptera checklist 2011

<i>Actidium aterrimum</i> (Motschulsky, 1845)	0	0	0	1	0	1
<i>Oligella foveolata</i> (Allibert, 1844)	1	0	10	2	0	13
<i>Oligella intermedia</i> Besuchet, 1971	0	0	1	0	0	1
<i>Ptilium horioni</i> Rosskothén, 1934	0	0	4	0	0	4
<i>Ptilium myrmecophilum</i> (Allibert, 1844)	0	5	3	0	0	8
<i>Millidium minutissimum</i> (Ljungh, 1804)	0	1	3	0	1	5
<i>Ptinella aptera</i> (Guérin-Méneville, 1839)	0	1	2	0	0	3
<i>Ptinella britannica</i> Matthews, A., 1858	0	1	0	1	0	2
<i>Ptinella cavelli</i> (Broun, 1893)	2	3	9	3	0	17
<i>Ptinella denticollis</i> (Fairmaire, 1858)	0	2	3	2	0	7
<i>Ptinella errabunda</i> Johnson, 1975	0	2	14	3	1	20
<i>Pteryx suturalis</i> (Heer, 1841)	0	3	8	6	2	19
<i>Nephanes titan</i> (Newman, 1834)	12	5	22	7	7	53
<i>Smicrus filicornis</i> (Fairmaire & Laboulbène, 1855)	0	2	0	0	0	2
<i>Baeocrara variolosa</i> (Mulsant & Rey, 1861)	0	0	0	1	0	1
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>arnoldi</i> Rosskothén, 1935	1	0	0	0	0	1
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>atomaria</i> (De Geer, 1774)	11	12	50	20	4	97
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>cephalotes</i> (Allibert, 1844)	0	0	1	0	0	1
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>cognata</i> (Matthews, A., 1877)	23	15	115	31	18	202
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>danica</i> Sundt, 1958	7	3	30	14	3	57
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>fascicularis</i> (Herbst, 1793)	30	25	122	42	10	229
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>henrici</i> (Matthews, A., 1872)	10	0	23	2	0	35
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>insularis</i> (Mäklin, 1852)	6	8	20	4	0	38
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>intermedia</i> (Gillmeister, 1845)	8	12	82	27	3	132
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>josephi</i> (Matthews, A., 1872)	3	0	0	0	0	3
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>lucidula</i> Rosskothén, 1935	0	1	3	0	0	4
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>montandonii</i> (Allibert, 1844)	6	12	22	7	0	47
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>norvegica</i> Strand, A., 1941	0	0	2	2	0	4
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>parva</i> Rosskothén, 1935	1	0	2	1	0	4
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>pumila</i> (Erichson, 1845)	1	0	1	0	0	2
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>rosskotheni</i> Sundt, 1971	0	0	17	6	0	23
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>rugulosa</i> Rosskothén, 1935	0	2	2	1	0	5
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>sericans</i> (Heer, 1841)	11	6	37	11	3	68
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>silvatica</i> Rosskothén, 1935	1	1	8	0	0	10
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>sitkaensis</i> (Motschulsky, 1845)	20	2	45	25	4	96
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>strandii</i> Sundt, 1958	4	1	7	6	1	19
<i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>thoracica</i> (Waltl, 1838)	2	7	15	9	3	36
<i>Acrotrichis</i> (<i>Ctenopteryx</i>) <i>grandicollis</i> (Mannerheim, 1844)	22	19	59	17	6	123
<i>Acrotrichis</i> (<i>Ctenopteryx</i>) <i>sanctae-helenae</i> Johnson, 1972	0	1	1	0	0	2
<u>Leiodidae [93/77]</u>						
<i>Sogda suturalis</i> (Zetterstedt, 1828)	0	3	0	0	0	3
<i>Hydnobius latifrons</i> (Curtis, 1840)	0	0	1	0	0	1
<i>Hydnobius punctatus</i> (Sturm, 1807)	0	3	4	0	0	7
<i>Hydnobius spinipes</i> (Gyllenhal, 1813)	0	0	2	0	0	2
<i>Triarthron maerkelii</i> Märkel, 1840	0	2	0	0	0	2
<i>Leiodes badia</i> (Sturm, 1807)	0	2	2	0	0	4
<i>Leiodes calcarata</i> (Erichson, 1845)	3	8	29	7	3	50
<i>Leiodes ciliaris</i> (Schmidt, W.L.E., 1841)	0	1	0	0	0	1
<i>Leiodes cinnamomea</i> (Panzer, 1793)	1	2	2	0	0	5

Yorkshire Coleoptera checklist 2011

<i>Leiodes ferruginea</i> (Fabricius, 1787)	2	3	1	1	1	8
<i>Leiodes furva</i> (Erichson, 1845)	1	0	0	0	0	1
<i>Leiodes gallica</i> (Reitter, 1884)	0	1	0	0	0	1
<i>Leiodes gyllenhalii</i> Stephens, 1829	0	2	0	0	0	2
<i>Leiodes litura</i> Stephens, 1832	1	3	3	1	0	8
<i>Leiodes lucens</i> (Fairmaire, 1855)	0	0	3	0	0	3
<i>Leiodes lunicollis</i> (Rye, 1872)	1	1	0	0	0	2
<i>Leiodes nigrita</i> (Schmidt, W.L.E., 1841)	0	1	0	0	0	1
<i>Leiodes obesa</i> (Schmidt, W.L.E., 1841)	1	0	10	5	0	16
<i>Leiodes picea</i> (Panzer, 1797)	0	1	0	0	0	1
<i>Leiodes rufipennis</i> (Paykull, 1798)	0	0	4	0	1	5
<i>Leiodes rugosa</i> Stephens, 1829	0	2	0	0	0	2
<i>Leiodes strigipennis</i> Daffner, 1983	0	0	3	2	0	5
<i>Leiodes triepkii</i> (Schmidt, W.L.E., 1841)	0	0	2	0	0	2
<i>Liocyrtusa minuta</i> (Ahrens, 1812)	1	1	0	1	0	3
<i>Liocyrtusa vittata</i> (Curtis, 1840)	1	1	2	1	2	7
<i>Agaricophagus cephalotes</i> Schmidt, W.L.E., 1841	0	0	2	0	0	2
<i>Colenis immunda</i> (Sturm, 1807)	0	2	6	3	0	11
<i>Anisotoma glabra</i> (Fabricius, 1792)	0	0	1	0	0	1
<i>Anisotoma humeralis</i> (Fabricius, 1792)	5	9	67	10	4	95
<i>Anisotoma orbicularis</i> (Herbst, 1792)	0	5	8	1	0	14
<i>Amphicyllis globus</i> (Fabricius, 1792)	0	3	10	1	0	14
<i>Agathidium</i> (Cyphocele) <i>arcticum</i> Thomson, C.G., 1862	0	0	1	0	0	1
<i>Agathidium</i> (Cyphocele) <i>nigrinum</i> Sturm, 1807	2	3	18	4	0	27
<i>Agathidium</i> (Neoceleble) <i>convexum</i> Sharp, 1866	2	0	0	1	0	3
<i>Agathidium</i> (Neoceleble) <i>marginatum</i> Sturm, 1807	2	1	0	0	0	3
<i>Agathidium</i> (Neoceleble) <i>nigripenne</i> (Fabricius, 1792)	2	5	29	14	3	53
<i>Agathidium</i> (Neoceleble) <i>rotundatum</i> (Gyllenhal, 1827)	0	5	4	0	0	9
<i>Agathidium</i> (Neoceleble) <i>varians</i> Beck, 1817	10	6	32	5	1	54
<i>Agathidium</i> (Agathidium) <i>atrum</i> (Paykull, 1798)	1	4	19	3	0	27
<i>Agathidium</i> (Agathidium) <i>laevigatum</i> Erichson, 1845	18	3	23	4	1	49
<i>Agathidium</i> (Agathidium) <i>pisanum</i> Brisout, 1872	0	0	0	1	0	1
<i>Agathidium</i> (Agathidium) <i>seminulum</i> (Linnaeus, 1758)	0	2	3	5	0	10
<i>Colon</i> (Eurycolon) <i>latum</i> Kraatz, 1850	0	3	2	0	0	5
<i>Colon</i> (Myloechus) <i>appendiculatum</i> (Sahlberg, C.R., 1822)	0	1	0	0	0	1
<i>Colon</i> (Myloechus) <i>brunneum</i> (Latreille, 1807)	1	2	3	3	0	9
<i>Colon</i> (Myloechus) <i>dentipes</i> (Sahlberg, C.R., 1822)	0	1	0	1	0	2
<i>Colon</i> (Colon) <i>serripes</i> (Sahlberg, C.R., 1822)	0	2	1	0	0	3
<i>Nemadus colonoides</i> (Kraatz, 1851)	1	1	0	1	0	3
<i>Nargus</i> (Nargus) <i>velox</i> (Spence, 1813)	21	6	95	17	6	145
<i>Nargus</i> (Nargus) <i>wilkinii</i> (Spence, 1813)	3	6	59	5	1	74
<i>Nargus</i> (Demochrus) <i>anisotomoides</i> (Spence, 1813)	1	4	16	0	0	21
<i>Choleva</i> (Cholevopsis) <i>spadicea</i> (Sturm, 1839)	0	2	7	0	0	9
<i>Choleva</i> (Choleva) <i>agilis</i> (Illiger, 1798)	2	3	9	0	0	14
<i>Choleva</i> (Choleva) <i>angustata</i> (Fabricius, 1781)	7	2	12	3	0	24
<i>Choleva</i> (Choleva) <i>cisteloides</i> (Frölich, 1799)	0	0	1	0	0	1
<i>Choleva</i> (Choleva) <i>fagniezi</i> Jeannel, 1922	0	0	10	5	0	15
<i>Choleva</i> (Choleva) <i>glauca</i> Britten, 1918	1	0	2	4	0	7
<i>Choleva</i> (Choleva) <i>jeanneli</i> Britten, 1922	7	2	10	5	0	24
<i>Choleva</i> (Choleva) <i>oblonga</i> Latreille, 1807	1	1	6	4	0	12

Yorkshire Coleoptera checklist 2011

<i>Sciodrepoides fumatus</i> (Spence, 1813)	3	6	18	11	2	40
<i>Sciodrepoides watsoni</i> (Spence, 1813)	17	14	70	19	3	123
<i>Catops chrysomeloides</i> (Panzer, 1798)	9	4	10	8	0	31
<i>Catops coracinus</i> Kellner, 1846	5	5	15	6	3	34
<i>Catops fuliginosus</i> Erichson, 1837	15	4	66	11	0	96
<i>Catops fuscus</i> (Panzer, 1794)	3	1	4	1	0	9
<i>Catops grandicollis</i> Erichson, 1837	11	4	32	8	4	59
<i>Catops kirbii</i> (Spence, 1813)	6	7	21	5	2	41
<i>Catops longulus</i> Kellner, 1846	0	5	7	3	1	16
<i>Catops morio</i> (Fabricius, 1787)	20	8	32	11	2	73
<i>Catops nigricans</i> (Spence, 1813)	17	3	35	6	1	62
<i>Catops nigrita</i> Erichson, 1837	0	5	43	3	2	53
<i>Catops tristis</i> (Panzer, 1793)	11	12	97	15	2	137
<i>Catopidius depressus</i> (Murray, 1856)	1	0	1	0	0	2
<i>Ptomaphagus medius</i> Rey, 1889	3	3	10	3	1	20
<i>Ptomaphagus subvillosus</i> (Goeze, 1777)	14	19	109	33	3	178
<i>Ptomaphagus varicornis</i> (Rosenhauer, 1847)	1	0	0	0	0	1
<i>Leptinus testaceus</i> Müller, P.W.J., 1817	1	5	8	3	0	17

Scymaenidae [32/18]

<i>Eutheia schaumii</i> Kiesenwetter, 1858	0	2	1	0	0	3
<i>Eutheia scydmaenoides</i> Stephens, 1830	0	2	2	1	0	5
<i>Cephennium gallicum</i> Ganglbauer, 1899	2	0	5	0	0	7
<i>Euconnus</i> (<i>Cladoconnus</i>) <i>denticornis</i> (Müller, P.W.J. & Kunze, 1822)	0	2	0	0	0	2
<i>Euconnus</i> (<i>Euconnus</i>) <i>fimetarius</i> (Chaudoir, 1845)	0	1	1	0	0	2
<i>Euconnus</i> (<i>Euconnus</i>) <i>hirticollis</i> (Illiger, 1798)	1	2	1	8	0	12
<i>Euconnus</i> (<i>Napochus</i>) <i>duboisii</i> Méquignon, 1929	2	0	1	0	0	3
<i>Microscydmus nanus</i> (Schaum, 1844)	0	1	0	1	0	2
<i>Neuraphes</i> (<i>Neuraphes</i>) <i>angulatus</i> (Müller, P.W.J. & Kunze, 1822)	0	3	1	0	0	4
<i>Neuraphes</i> (<i>Neuraphes</i>) <i>elongatulus</i> (Müller, P.W.J. & Kunze, 1822)	0	9	27	7	1	44
<i>Neuraphes</i> (<i>Neuraphes</i>) <i>talparum</i> Lokay, 1921	0	3	0	1	0	4
<i>Neuraphes</i> (<i>Pararaphes</i>) <i>plicicollis</i> Reitter, 1879	0	0	0	3	0	3
<i>Scydmorephes helvolus</i> (Schaum, 1844)	4	0	5	1	0	10
<i>Scydmorephes sparshalli</i> (Denny, 1825)	0	2	0	1	0	3
<i>Stenichnus bicolor</i> (Denny, 1825)	0	2	8	3	0	13
<i>Stenichnus collaris</i> (Müller, P.W.J. & Kunze, 1822)	8	23	42	18	2	93
<i>Stenichnus scutellaris</i> (Müller, P.W.J. & Kunze, 1822)	8	4	9	1	0	22
<i>Scydmaenus</i> (<i>Scydmaenus</i>) <i>tarsatus</i> Müller, P.W.J. & Kunze, 1822	19	10	91	18	2	140

Silphidae [21/19]

<i>Necrodes littoralis</i> (Linnaeus, 1758)	6	5	19	9	28	67
<i>Thanatophilus dispar</i> (Herbst, 1793)	0	1	0	2	0	3
<i>Thanatophilus rugosus</i> (Linnaeus, 1758)	27	6	47	22	9	111
<i>Thanatophilus sinuatus</i> (Fabricius, 1775)	9	4	21	6	1	41
<i>Oiceoptoma thoracicum</i> (Linnaeus, 1758)	9	25	32	8	0	74
<i>Aclypea opaca</i> (Linnaeus, 1758)	5	7	6	6	0	24
<i>Aclypea undata</i> (Müller, O.F., 1776)	0	1	0	0	0	1
<i>Dendroxena quadrimaculata</i> (Scopoli, 1772)	3	2	22	2	0	29
<i>Silpha atrata</i> Linnaeus, 1758	12	27	49	15	7	110

Yorkshire Coleoptera checklist 2011

<i>Silpha laevigata</i> Fabricius, 1775	0	2	3	0	0	5
<i>Silpha obscura</i> Linnaeus, 1758	0	0	1	0	0	1
<i>Silpha tristis</i> Illiger, 1798	0	1	42	7	1	51
<i>Silpha tyrolensis</i> Laicharting, 1781	0	3	5	2	3	13
<i>Nicrophorus humator</i> (Gleditsch, 1767)	22	3	87	40	127	279
<i>Nicrophorus interruptus</i> Stephens, 1830	0	1	3	0	0	4
<i>Nicrophorus investigator</i> Zetterstedt, 1824	12	4	33	28	105	182
<i>Nicrophorus vespillo</i> (Linnaeus, 1758)	10	7	36	15	4	72
<i>Nicrophorus vespilloides</i> Herbst, 1783	6	20	123	6	3	158
<i>Nicrophorus vestigator</i> Herschel, 1807	0	4	0	0	0	4

Staphylinidae [1080/802]

<i>Acidota crenata</i> (Fabricius, 1792)	2	12	12	10	4	40
<i>Acidota cruentata</i> Mannerheim, 1830	0	3	8	7	0	18
<i>Anthobium atrocephalum</i> (Gyllenhal, 1827)	14	16	156	18	1	205
<i>Anthobium unicolor</i> (Marsham, 1802)	34	25	194	48	7	308
<i>Anthophagus</i> (<i>Anthophagus</i>) <i>alpinus</i> (Paykull, 1790)	0	0	0	4	0	4
<i>Anthophagus</i> (<i>Phaganthus</i>) <i>caraboides</i> (Linnaeus, 1758)	0	27	4	9	4	44
<i>Deliphrum tectum</i> (Paykull, 1789)	2	3	16	4	2	27
<i>Eucnecosum brachypterum</i> (Gravenhorst, 1802)	0	1	6	8	5	20
<i>Geodromicus nigrita</i> (Müller, P.W.J., 1821)	0	3	4	7	4	18
<i>Lesteva hanseni</i> Lohse, 1953	0	0	0	3	0	3
<i>Lesteva longoelytrata</i> (Goeze, 1777)	43	43	125	48	20	279
<i>Lesteva monticola</i> Kiesenwetter, 1847	0	4	2	8	10	24
<i>Lesteva pubescens</i> Mannerheim, 1830	1	11	12	5	2	31
<i>Lesteva punctata</i> Erichson, 1839	3	9	11	6	2	31
<i>Lesteva sicula</i> subsp. <i>heeri</i> Fauvel, 1871	37	40	166	55	13	311
<i>Olophrum consimile</i> (Gyllenhal, 1810)	0	2	0	0	0	2
<i>Olophrum fuscum</i> (Gravenhorst, 1806)	2	3	20	8	6	39
<i>Olophrum piceum</i> (Gyllenhal, 1810)	20	33	123	26	10	212
<i>Philorinum sordidum</i> (Stephens, 1834)	13	13	25	6	4	61
<i>Phyllodrepoidea crenata</i> Ganglbauer, 1895	0	8	11	3	0	22
<i>Eusphalerum luteum</i> (Marsham, 1802)	1	8	18	6	1	34
<i>Eusphalerum minutum</i> (Fabricius, 1792)	8	10	4	36	7	65
<i>Eusphalerum primulae</i> (Stephens, 1834)	3	12	14	4	0	33
<i>Eusphalerum sorbi</i> (Gyllenhal, 1810)	4	20	12	16	4	56
<i>Eusphalerum torquatum</i> (Marsham, 1802)	3	4	14	11	8	40
<i>Acrolocha minuta</i> (Olivier, 1795)	0	1	2	0	0	3
<i>Acrolocha sulcula</i> (Stephens, 1834)	5	8	11	4	2	30
<i>Acrulia inflata</i> (Gyllenhal, 1813)	3	6	43	8	0	60
<i>Dropephylla devillei</i> Bernhauer, 1902	0	6	7	4	0	17
<i>Dropephylla ioptera</i> (Stephens, 1832)	6	4	8	7	1	26
<i>Dropephylla vilis</i> (Erichson, 1840)	3	11	43	9	0	66
<i>Hapalareae pygmaea</i> (Paykull, 1800)	0	1	7	3	0	11
<i>Micralymma marinum</i> (Ström, 1783)	4	9	1	0	0	14
<i>Omalius</i> <i>Gravenhorst</i> , 1802	0	0	1	0	0	1
<i>Omalius allardi</i> Fairmaire & Brisout, 1859	0	3	4	4	0	11
<i>Omalius caesum</i> Gravenhorst, 1806	36	9	86	16	2	149
<i>Omalius excavatum</i> Stephens, 1834	26	8	33	15	2	84
<i>Omalius exiguum</i> Gyllenhal, 1810	0	1	2	0	0	3

Yorkshire Coleoptera checklist 2011

<i>Omalium italicum</i> Bernhauer, 1902	7	2	32	4	0	45
<i>Omalium laeviusculum</i> Gyllenhal, 1827	5	6	0	0	0	11
<i>Omalium laticolle</i> Kraatz, 1858	0	0	3	4	0	7
<i>Omalium oxyacanthae</i> Gravenhorst, 1806	2	4	9	6	1	22
<i>Omalium riparium</i> Thomson, C.G., 1857	10	3	0	0	0	13
<i>Omalium rivulare</i> (Paykull, 1789)	39	19	181	34	7	280
<i>Omalium rugatum</i> Mulsant & Rey, 1880	0	1	7	3	0	11
<i>Omalium rugulipenne</i> Rye, 1864	0	2	0	0	0	2
<i>Omalium septentrionis</i> Thomson, C.G., 1856	0	6	8	3	0	17
<i>Phloeonomus punctipennis</i> Thomson, C.G., 1867	3	6	24	12	3	48
<i>Phloeonomus pusillus</i> (Gravenhorst, 1806)	3	5	20	2	1	31
<i>Phloeostiba plana</i> (Paykull, 1792)	0	2	0	0	0	2
<i>Phyllodrepa floralis</i> (Paykull, 1789)	13	6	11	5	0	35
<i>Phyllodrepa puberula</i> Bernhauer, 1903	0	1	1	0	0	2
<i>Phyllodrepa salicis</i> (Gyllenhal, 1810)	0	1	0	0	0	1
<i>Xylodromus concinnus</i> (Marsham, 1802)	14	16	16	8	2	56
<i>Xylodromus depressus</i> (Gravenhorst, 1802)	5	3	7	6	1	22
<i>Xylostiba monilicornis</i> (Gyllenhal, 1810)	0	0	3	0	0	3
<i>Coryphium angusticolle</i> Stephens, 1834	3	3	27	6	0	39
<i>Euedectus whitei</i> Sharp, 1871	0	0	0	3	0	3
<i>Megarathrus bellevoeyi</i> (Saulcy, 1862)	4	5	19	0	0	28
<i>Megarathrus denticollis</i> (Beck, 1817)	11	11	80	21	6	129
<i>Megarathrus depressus</i> (Paykull, 1789)	21	19	138	37	3	218
<i>Megarathrus prosseni</i> Schatzmayr, 1904	1	0	7	0	0	8
<i>Metopsia clypeata</i> (Müller, P.W.J., 1821)	24	10	56	28	0	118
<i>Proteinus atomarius</i> Erichson, 1840	0	1	6	2	0	9
<i>Proteinus brachypterus</i> (Fabricius, 1792)	10	24	90	22	3	149
<i>Proteinus crenulatus</i> Pandellé, 1867	0	3	12	2	0	17
<i>Proteinus laevigatus</i> Hochhuth, 1871	6	1	31	9	1	48
<i>Proteinus ovalis</i> Stephens, 1834	39	14	190	32	3	278
<i>Arrhenopeplus tesserula</i> Curtis, 1828	0	0	2	3	1	6
<i>Micropeplus fulvus</i> Erichson, 1840	14	6	46	18	0	84
<i>Micropeplus porcatus</i> (Paykull, 1789)	2	10	1	4	0	17
<i>Micropeplus staphylinoides</i> (Marsham, 1802)	17	7	33	6	1	64
<i>Trichonyx sulcicollis</i> (Reichenbach, 1816)	0	1	1	0	0	2
<i>Trimium brevicorne</i> (Reichenbach, 1816)	0	2	1	1	0	4
<i>Bibloporus bicolor</i> (Denny, 1825)	0	2	7	0	0	9
<i>Bibloporus minutus</i> Raffray, 1914	0	3	1	1	0	5
<i>Biblopectus ambiguus</i> (Reichenbach, 1816)	1	1	1	4	0	7
<i>Biblopectus pusillus</i> (Denny, 1825)	0	0	0	1	0	1
<i>Biblopectus tenebrosus</i> (Reitter, 1880)	0	0	0	3	0	3
<i>Euplectus bescidicus</i> Reitter, 1881	0	1	0	0	0	1
<i>Euplectus karstenii</i> (Reichenbach, 1816)	0	0	3	3	0	6
<i>Euplectus nanus</i> (Reichenbach, 1816)	3	1	3	1	1	9
<i>Euplectus piceus</i> Motschulsky, 1835	0	2	1	2	0	5
<i>Euplectus sanguineus</i> Denny, 1825	6	4	21	3	2	36
<i>Euplectus signatus</i> (Reichenbach, 1816)	0	2	10	0	0	12
<i>Plectophloeus nitidus</i> (Fairmaire, 1857)	0	0	1	0	0	1
<i>Batrisodes venustus</i> (Reichenbach, 1816)	1	1	0	1	0	3
<i>Bryaxis bulbifer</i> (Reichenbach, 1816)	10	9	22	35	2	78

Yorkshire Coleoptera checklist 2011

<i>Bryaxis curtisii</i> (Leach, 1817)	0	4	5	7	0	16
<i>Bryaxis puncticollis</i> (Denny, 1825)	5	12	17	19	1	54
<i>Bythinus burrellii</i> Denny, 1825	2	10	7	15	2	36
<i>Bythinus macropalpus</i> Aubé, 1833	3	2	9	3	0	17
<i>Tychus niger</i> (Paykull, 1800)	14	19	22	13	3	71
<i>Brachygluta fossulata</i> (Reichenbach, 1816)	8	10	27	19	0	64
<i>Brachygluta helferi</i> (Schmidt-Göbel, 1836)	19	0	0	0	0	19
<i>Fagniezia impressus</i> (Panzer, 1803)	0	2	1	3	0	6
<i>Reichenbachia juncorum</i> (Leach, 1817)	5	4	3	5	0	17
<i>Rybaxis longicornis</i> (Leach, 1817)	11	0	46	8	1	66
<i>Pselaphaulax dresdensis</i> (Herbst, 1792)	2	0	1	2	0	5
<i>Pselaphus heisei</i> Herbst, 1792	6	9	9	14	3	41
<i>Claviger testaceus</i> Preyssler, 1790	0	3	0	0	0	3
<i>Phloeocharis subtilissima</i> Mannerheim, 1830	4	9	29	9	2	53
<i>Cilea silphoides</i> (Linnaeus, 1767)	8	9	16	5	0	38
<i>Lamprinodes saginatus</i> (Gravenhorst, 1806)	8	2	5	0	0	15
<i>Sepedophilus constans</i> (Fowler, 1888)	0	1	1	0	0	2
<i>Sepedophilus immaculatus</i> (Stephens, 1832)	14	2	16	1	0	33
<i>Sepedophilus littoreus</i> (Linnaeus, 1758)	3	6	21	2	0	32
<i>Sepedophilus lusitanicus</i> Hammond, 1973	0	0	2	0	0	2
<i>Sepedophilus marshami</i> (Stephens, 1832)	66	22	209	23	6	326
<i>Sepedophilus nigripennis</i> (Stephens, 1832)	76	20	200	27	7	330
<i>Sepedophilus pedicularius</i> (Gravenhorst, 1802)	1	0	8	10	0	19
<i>Sepedophilus testaceus</i> (Fabricius, 1792)	2	0	0	2	0	4
<i>Tachinus corticinus</i> Gravenhorst, 1802	8	10	21	36	4	79
<i>Tachinus elongatus</i> Gyllenhal, 1810	0	15	8	4	4	31
<i>Tachinus humeralis</i> Gravenhorst, 1802	9	8	46	7	2	72
<i>Tachinus laticollis</i> Gravenhorst, 1802	14	18	100	13	4	149
<i>Tachinus lignorum</i> (Linnaeus, 1758)	0	1	3	3	0	7
<i>Tachinus marginellus</i> (Fabricius, 1781)	33	27	207	54	17	338
<i>Tachinus pallipes</i> (Gravenhorst, 1806)	4	10	50	8	1	73
<i>Tachinus proximus</i> Kraatz, 1855	2	5	20	7	1	35
<i>Tachinus rufipennis</i> Gyllenhal, 1810	0	1	0	1	0	2
<i>Tachinus rufipes</i> (Linnaeus, 1758)	73	57	471	108	17	726
<i>Tachinus subterraneus</i> (Linnaeus, 1758)	9	4	33	8	1	55
<i>Tachyporus</i> Gravenhorst, 1802	0	0	1	0	0	1
<i>Tachyporus atriceps</i> Stephens, 1832	9	7	11	6	0	33
<i>Tachyporus chrysomelinus</i> (Linnaeus, 1758)	20	3	218	26	3	270
<i>Tachyporus dispar</i> (Paykull, 1789)	65	68	338	73	15	559
<i>Tachyporus formosus</i> Matthews, A.H., 1838	0	1	0	1	1	3
<i>Tachyporus hypnorum</i> (Fabricius, 1775)	139	85	584	98	18	924
<i>Tachyporus nitidulus</i> (Fabricius, 1781)	52	39	237	38	7	373
<i>Tachyporus obtusus</i> (Linnaeus, 1767)	81	51	449	63	15	659
<i>Tachyporus pallidus</i> Sharp, 1871	38	24	116	25	2	205
<i>Tachyporus pusillus</i> Gravenhorst, 1806	8	4	32	9	3	56
<i>Tachyporus quadriscopulatus</i> Pandellé, 1869	0	0	0	1	0	1
<i>Tachyporus solutus</i> Erichson, 1839	22	8	256	2	1	289
<i>Tachyporus tersus</i> Erichson, 1839	0	10	8	5	1	24
<i>Tachyporus transversalis</i> Gravenhorst, 1806	6	6	28	4	0	44
<i>Bolitobius cingulatus</i> (Mannerheim, 1830)	5	8	28	6	1	48

Yorkshire Coleoptera checklist 2011

<i>Ischnosoma splendidum</i> (Gravenhorst, 1806)	0	0	8	2	0	10
<i>Lordithon exoletus</i> (Erichson, 1839)	2	5	9	1	0	17
<i>Lordithon lunulatus</i> (Linnaeus, 1761)	4	12	34	10	5	65
<i>Lordithon thoracicus</i> (Fabricius, 1777)	4	12	58	18	2	94
<i>Lordithon trinotatus</i> (Erichson, 1839)	8	16	73	11	3	111
<i>Mycetoporus angularis</i> Mulsant & Rey, 1853	0	1	1	0	0	2
<i>Mycetoporus baudueri</i> Mulsant & Rey, 1875	0	1	0	0	0	1
<i>Mycetoporus clavicornis</i> (Stephens, 1832)	3	14	23	5	2	47
<i>Mycetoporus despectus</i> Strand, A., 1969	0	0	4	1	1	6
<i>Mycetoporus lepidus</i> (Gravenhorst, 1806)	8	25	45	21	3	102
<i>Mycetoporus longulus</i> Mannerheim, 1830	4	8	10	10	2	34
<i>Mycetoporus nigricollis</i> Stephens, 1835	20	16	38	14	1	89
<i>Mycetoporus piceolus</i> Rey, 1883	2	1	0	0	0	3
<i>Mycetoporus punctus</i> (Gravenhorst, 1806)	2	3	4	0	0	9
<i>Mycetoporus rufescens</i> (Stephens, 1832)	2	12	15	4	2	35
<i>Parabolitobius inclinans</i> (Gravenhorst, 1806)	0	9	6	4	0	19
<i>Trichophya pilicornis</i> (Gyllenhal, 1810)	0	3	4	0	0	7
<i>Habrocerus capillaricornis</i> (Gravenhorst, 1806)	33	13	108	40	8	202
<i>Deinopsis erosa</i> (Stephens, 1832)	4	0	4	2	0	10
<i>Gymnusa brevicollis</i> (Paykull, 1800)	3	9	9	2	0	23
<i>Gymnusa variegata</i> Kiesenwetter, 1845	2	3	7	0	2	14
<i>Cypha discoidea</i> (Erichson, 1839)	3	0	0	0	0	3
<i>Cypha laeviuscula</i> (Mannerheim, 1830)	15	8	22	5	1	51
<i>Cypha longicornis</i> (Paykull, 1800)	33	24	96	9	2	164
<i>Cypha pulicaria</i> (Erichson, 1839)	4	3	5	1	0	13
<i>Cypha punctum</i> (Motschulsky, 1857)	0	2	0	0	0	2
<i>Cypha seminulum</i> (Erichson, 1839)	0	3	0	1	0	4
<i>Oligota apicata</i> (Erichson, 1837)	5	3	5	2	2	17
<i>Holobus flavicornis</i> (Lacordaire, 1835)	0	0	1	0	0	1
<i>Oligota granaria</i> Erichson, 1837	1	0	0	0	0	1
<i>Oligota inflata</i> (Mannerheim, 1830)	11	13	19	2	0	45
<i>Oligota parva</i> Kraatz, 1862	8	1	14	1	0	24
<i>Oligota picipes</i> (Stephens, 1832)	4	2	3	0	0	9
<i>Oligota pumilio</i> Kiesenwetter, 1858	1	0	1	1	0	3
<i>Oligota punctulata</i> Heer, 1839	4	1	39	8	4	56
<i>Oligota pusillima</i> (Gravenhorst, 1806)	9	3	9	1	0	22
<i>Myllaena brevicornis</i> (Matthews, A.H., 1838)	7	13	39	15	12	86
<i>Myllaena dubia</i> (Gravenhorst, 1806)	15	4	22	2	4	47
<i>Myllaena elongata</i> (Matthews, A.H., 1838)	3	2	1	0	0	6
<i>Myllaena gracilis</i> (Matthews, A.H., 1838)	3	0	0	2	0	5
<i>Myllaena graeca</i> Kraatz, 1858	1	0	0	0	0	1
<i>Myllaena infuscata</i> Kraatz, 1853	8	2	8	9	1	28
<i>Myllaena intermedia</i> Erichson, 1837	3	2	5	1	0	11
<i>Myllaena kraatzi</i> Sharp, 1871	2	4	2	0	1	9
<i>Myllaena minuta</i> (Gravenhorst, 1806)	6	1	7	10	2	26
<i>Diglotta mersa</i> (Haliday, 1837)	3	0	0	0	0	3
<i>Hygronoma dimidiata</i> (Gravenhorst, 1806)	21	5	41	12	3	82
<i>Encephalus complicans</i> Stephens, 1832	5	5	22	8	1	41
<i>Gyrophaena affinis</i> Mannerheim, 1830	4	5	16	5	0	30
<i>Gyrophaena manca</i> Erichson, 1839	3	9	32	10	1	55

Yorkshire Coleoptera checklist 2011

<i>Gyrophana bihamata</i> Thomson,C.G.,1867	2	0	8	2	0	12
<i>Gyrophana congrua</i> Erichson,1837	1	2	1	0	0	4
<i>Gyrophana fasciata</i> (Marsham,1802)	3	8	21	12	1	45
<i>Gyrophana gentilis</i> Erichson,1839	4	13	41	11	3	72
<i>Gyrophana joyi</i> Wendeler,1924	0	0	0	1	1	2
<i>Gyrophana joyioides</i> Wusthoff,1937	0	0	7	5	0	12
<i>Agaricochara latissima</i> (Stephens,1832)	2	5	43	7	2	59
<i>Gyrophana lucidula</i> Erichson,1837	1	1	0	0	0	2
<i>Gyrophana minima</i> Erichson,1837	1	4	4	6	0	15
<i>Gyrophana munsteri</i> Strand,A.,1935	0	1	0	1	0	2
<i>Gyrophana nana</i> (Paykull,1800)	2	0	1	1	1	5
<i>Gyrophana poweri</i> Crotch,1867	1	0	0	0	0	1
<i>Gyrophana strictula</i> Erichson,1839	1	0	4	0	0	5
<i>Gyrophana williamsi</i> Strand,A.,1935	0	0	1	1	0	2
<i>Placusa depressa</i> Maklin,1845	0	1	3	0	0	4
<i>Placusa tachyporoides</i> (Waltl,1838)	0	0	0	1	0	1
<i>Homalota plana</i> (Gyllenhal,1810)	2	7	15	7	1	32
<i>Anomognathus cuspidatus</i> (Erichson,1839)	1	4	8	0	1	14
<i>Phytosus balticus</i> Kraatz,1859	4	2	0	0	0	6
<i>Phytosus spinifer</i> Curtis,1838	3	1	0	0	0	4
<i>Arena tabida</i> (Kiesenwetter,1850)	2	0	0	0	0	2
<i>Thecturota marchii</i> (Dodero,1922)	0	0	1	0	0	1
<i>Leptusa fumida</i> (Erichson,1839)	11	45	217	21	6	300
<i>Leptusa norvegica</i> Strand,A.,1941	0	3	1	0	0	4
<i>Leptusa pulchella</i> (Mannerheim,1830)	0	9	15	4	2	30
<i>Leptusa ruficollis</i> (Erichson,1839)	3	19	31	8	1	62
<i>Heterota plumbea</i> (Waterhouse,G.R.,1858)	2	0	0	0	0	2
<i>Bolitochara bella</i> Markel,1844	2	0	1	1	0	4
<i>Bolitochara lucida</i> (Gravenhorst,1802)	2	5	16	7	0	30
<i>Bolitochara mulsanti</i> Sharp,1875	0	3	5	0	1	9
<i>Bolitochara obliqua</i> Erichson,1837	14	23	139	23	6	205
<i>Bolitochara pulchra</i> (Gravenhorst,1806)	0	0	1	0	0	1
<i>Autalia impressa</i> (Olivier,1795)	4	12	116	16	3	151
<i>Autalia longicornis</i> Scheerpeltz,1947	1	2	22	7	1	33
<i>Autalia puncticollis</i> Sharp,1864	0	0	0	2	1	3
<i>Autalia rivularis</i> (Gravenhorst,1802)	23	15	115	26	7	186
<i>Cordalia obscura</i> (Gravenhorst,1802)	0	3	1	2	0	6
<i>Falagria caesa</i> Erichson,1837	1	4	16	1	0	22
<i>Falagria sulcatula</i> (Gravenhorst,1806)	0	0	2	0	0	2
<i>Falagrioma thoracica</i> (Stephens,1832)	0	1	2	0	0	3
<i>Myrmecopora sulcata</i> (Kiesenwetter,1850)	3	2	0	0	0	5
<i>Myrmecopora uvida</i> (Erichson,1840)	0	1	0	0	0	1
<i>Thinonoma atra</i> (Gravenhorst,1806)	31	3	38	8	1	81
<i>Tachyusa coarctata</i> (Erichson,1837)	0	1	1	0	0	2
<i>Tachyusa constricta</i> (Erichson,1837)	3	11	1	5	0	20
<i>Ischnopoda leucopus</i> (Marsham,1802)	4	11	5	4	0	24
<i>Ischnopoda umbratica</i> (Erichson,1837)	0	2	0	0	0	2
<i>Gnypeta caerulea</i> (Sahlberg,C.R.,1831)	0	1	0	4	4	9
<i>Gnypeta carbonaria</i> (Mannerheim,1830)	11	8	26	5	0	50
<i>Gnypeta ripicola</i> (Kiesenwetter,1844)	0	1	2	0	0	3

Yorkshire Coleoptera checklist 2011

<i>Gnypeta rubrior</i> Tottenham,1939	4	1	8	0	0	13
<i>Dasygnypeta velata</i> (Erichson,1837)	2	0	0	0	0	2
<i>Callicerus obscurus</i> Gravenhorst,1802	6	3	7	3	0	19
<i>Callicerus rigidicornis</i> (Erichson,1839)	1	1	15	3	1	21
<i>Schistoglossa aubei</i> (Brisout,1860)	1	0	0	0	0	1
<i>Schistoglossa curtipennis</i> (Sharp,1869)	1	0	7	0	0	8
<i>Schistoglossa gemina</i> (Erichson,1837)	3	0	3	2	0	8
<i>Boreophilia eremita</i> (Rye,1866)	4	8	27	8	3	50
<i>Dacrila fallax</i> (Kraatz,1856)	6	0	3	0	0	9
<i>Hydrosmeeta delicatula</i> (Sharp,1869)	1	0	0	0	0	1
<i>Hydrosmeeta eximia</i> (Sharp,1869)	1	1	0	1	0	3
<i>Hydrosmeeta fragilis</i> (Kraatz,1854)	1	1	0	0	0	2
<i>Hydrosmeeta subtilissima</i> (Kraatz,1854)	0	0	0	1	0	1
<i>Aloconota cambrica</i> (Wollaston,1855)	0	3	0	2	0	5
<i>Aloconota currax</i> (Kraatz,1856)	2	11	3	1	2	19
<i>Aloconota gregaria</i> (Erichson,1839)	83	50	139	39	7	318
<i>Aloconota insecta</i> (Thomson,C.G.,1856)	2	3	13	3	2	23
<i>Aloconota sulcifrons</i> (Stephens,1832)	5	1	22	1	2	31
<i>Aloconota languida</i> (Erichson,1837)	3	0	0	1	1	5
<i>Pycnota paradoxa</i> (Mulsant & Rey,1861)	3	0	0	1	0	4
<i>Amischa analis</i> (Gravenhorst,1802)	39	59	194	34	10	336
<i>Amischa bifoveolata</i> (Mannerheim,1830)	21	11	18	6	3	59
<i>Amischa decipiens</i> (Sharp,1869)	20	6	38	7	2	73
<i>Amischa forcipata</i> Mulsant & Rey,1873	4	0	5	2	0	11
<i>Amischa nigrofusca</i> (Stephens,1829)	3	1	11	2	2	19
<i>Amidobia talpa</i> (Heer,1841)	2	7	6	0	0	15
<i>Nehemitropia lividipennis</i> (Mannerheim,1830)	22	11	53	14	0	100
<i>Notothecta confusa</i> (Markel,1844)	0	0	1	0	0	1
<i>Notothecta flavipes</i> (Gravenhorst,1806)	1	7	14	0	0	22
<i>Lyprocorrhe anceps</i> (Erichson,1837)	1	5	5	1	0	12
<i>Neohilara subterranea</i> (Mulsant & Rey,1853)	0	0	2	0	0	2
<i>Brundinia meridionalis</i> (Mulsant & Rey,1853)	10	0	0	0	0	10
<i>Dochmonota clancula</i> (Erichson,1837)	2	0	12	6	0	20
<i>Geostiba circellaris</i> (Gravenhorst,1806)	25	27	221	53	11	337
<i>Ousipalia caesula</i> (Erichson,1839)	0	1	0	0	0	1
<i>Dinaraea aequata</i> (Erichson,1837)	16	12	59	13	3	103
<i>Dinaraea angustula</i> (Gyllenhal,1810)	13	4	31	3	1	52
<i>Dinaraea linearis</i> (Gravenhorst,1802)	0	1	3	0	0	4
<i>Dadobia immersa</i> (Erichson,1837)	2	1	4	4	0	11
<i>Plataraea brunnea</i> (Fabricius,1798)	8	5	26	2	1	42
<i>Liogluta granigera</i> (Kiesenwetter,1850)	1	8	11	2	0	22
<i>Liogluta longiuscula</i> (Gravenhorst,1802)	9	4	9	2	1	25
<i>Liogluta alpestris</i> (Heer,1839)	4	46	9	1	1	61
<i>Liogluta microptera</i> Thomson,C.G.,1867	6	4	56	10	3	79
<i>Liogluta pagana</i> (Erichson,1839)	3	1	0	3	0	7
<i>Philhygra arctica</i> (Thomson,C.G.,1856)	0	6	1	1	0	8
<i>Philhygra britteni</i> Joy,1913	10	1	5	0	0	16
<i>Philhygra debilis</i> (Erichson,1837)	8	0	8	0	1	17
<i>Philhygra deformis</i> (Kraatz,1856)	1	0	0	0	1	2
<i>Philhygra elongatula</i> (Gravenhorst,1802)	30	31	39	23	9	132

Yorkshire Coleoptera checklist 2011

<i>Philhygra fallaciosa</i> (Sharp,1869)	1	2	1	0	0	4
<i>Philhygra gyllenhali</i> (Thomson,C.G.,1856)	7	2	10	2	0	21
<i>Philhygra hygrobica</i> (Thomson,C.G.,1856)	3	1	4	2	0	10
<i>Philhygra hygrotopora</i> (Kraatz,1856)	6	6	12	4	4	32
<i>Philhygra luridipennis</i> (Mannerheim,1830)	6	3	8	5	0	22
<i>Philhygra malleus</i> (Joy,1913)	27	3	30	5	2	67
<i>Philhygra melanocera</i> (Thomson,C.G.,1856)	10	4	44	4	0	62
<i>Philhygra obtusangula</i> (Joy,1913)	1	4	13	2	3	23
<i>Philhygra palustris</i> (Kiesenwetter,1844)	10	6	8	2	0	26
<i>Philhygra terminalis</i> (Gravenhorst,1806)	4	0	0	0	0	4
<i>Philhygra volans</i> (Scriba,1859)	5	0	12	1	1	19
<i>Parameotica difficilis</i> (Brisout,1860)	17	0	0	0	0	17
<i>Dilacra luteipes</i> (Erichson,1837)	5	1	3	4	0	13
<i>Dilacra vilis</i> (Erichson,1837)	1	0	4	6	0	11
<i>Enalodroma hepatica</i> (Erichson,1839)	1	2	5	6	2	16
<i>Bessobia excellens</i> (Kraatz,1856)	2	5	2	6	0	15
<i>Bessobia fungivora</i> (Thomson,C.G.,1867)	4	4	3	5	0	16
<i>Bessobia monticola</i> (Thomson,C.G.,1852)	1	2	6	3	0	12
<i>Bessobia occulta</i> (Erichson,1837)	0	3	3	0	1	7
<i>Atheta divisa</i> (Markel,1844)	1	0	0	1	0	2
<i>Atheta euryptera</i> (Stephens,1832)	2	2	2	1	0	7
<i>Atheta harwoodi</i> (Williams,1930)	6	4	77	11	5	103
<i>Atheta liturata</i> (Stephens,1832)	0	5	5	14	6	30
<i>Atheta vaga</i> (Heer,1839)	4	12	52	12	1	81
<i>Atheta nigritula</i> (Gravenhorst,1802)	2	0	5	0	0	7
<i>Anopleta corvina</i> (Thomson,C.G.,1856)	0	4	7	0	2	13
<i>Anopleta puberula</i> (Sharp,1869)	0	0	1	0	0	1
<i>Microdota amicula</i> (Stephens,1832)	32	20	95	30	9	186
<i>Microdota atricolor</i> (Sharp,1869)	5	4	19	4	2	34
<i>Microdota benickiella</i> (Brundin,1948)	0	0	1	2	0	3
<i>Microdota boreella</i> (Brundin,1948)	0	0	2	1	0	3
<i>Microdota excelsa</i> (Bernhauer,1911)	0	0	1	0	0	1
<i>Microdota indubia</i> (Sharp,1869)	10	6	39	15	3	73
<i>Microdota liliputana</i> (Brisout,1860)	0	0	2	0	0	2
<i>Microdota minuscula</i> (Brisout,1860)	0	0	1	0	0	1
<i>Microdota spatuloides</i> (Benick,1939)	0	1	0	0	0	1
<i>Microdota subtilis</i> (Scriba,1866)	0	0	2	2	0	4
<i>Cadaverota cadaverina</i> (Brisout,1860)	0	9	18	0	0	27
<i>Alaobia gagatina</i> (Baudi,1848)	3	12	16	1	1	33
<i>Alaobia hybrida</i> (Sharp,1869)	0	2	0	0	0	2
<i>Alaobia pallidicornis</i> (Thomson,C.G.,1856)	3	4	25	9	1	42
<i>Alaobia sodalis</i> (Erichson,1837)	1	12	16	6	3	38
<i>Alaobia taxiceroides</i> Munster,1932	0	1	5	0	0	6
<i>Oreostiba tibialis</i> (Heer,1839)	0	0	0	6	3	9
<i>Alaobia trinotata</i> (Kraatz,1856)	36	28	132	23	3	222
<i>Coprothassa melanaria</i> (Mannerheim,1830)	4	1	0	0	0	5
<i>Mocyta amplicollis</i> (Mulsant & Rey,1873)	13	5	23	3	1	45
<i>Mocyta clientula</i> (Erichson,1839)	3	4	5	3	0	15
<i>Mocyta fungi</i> (Gravenhorst,1806)	86	91	681	114	29	1001
<i>Mocyta gilvicollis</i> Scheerpeltz	0	2	1	0	0	3

Yorkshire Coleoptera checklist 2011

<i>Mocyta orbata</i> (Erichson,1837)	4	1	16	2	1	24
<i>Mocyta orphana</i> (Erichson,1837)	1	0	1	0	0	2
<i>Acrotona aterrima</i> (Gravenhorst,1802)	30	13	125	17	8	193
<i>Acrotona benicki</i> (Allen,1940)	0	1	0	0	0	1
<i>Acrotona exigua</i> (Erichson,1837)	0	0	1	0	0	1
<i>Acrotona muscorum</i> (Brisout,1860)	11	9	31	16	5	72
<i>Acrotona obfusca</i> (Gravenhorst,1802)	0	0	3	0	0	3
<i>Acrotona parvula</i> (Mannerheim,1830)	11	10	42	13	9	85
<i>Acrotona pygmaea</i> (Gravenhorst,1802)	2	0	31	9	0	42
<i>Rhagocneme subsinuata</i> (Erichson,1839)	1	2	10	1	0	14
<i>Datomicra canescens</i> (Sharp,1869)	0	0	3	0	0	3
<i>Datomicra celata</i> (Erichson,1837)	38	28	251	28	14	359
<i>Datomicra dadopora</i> (Thomson,C.G.,1867)	4	4	49	10	2	69
<i>Datomicra nigra</i> (Kraatz,1856)	46	16	131	38	9	240
<i>Datomicra sordidula</i> (Erichson,1837)	7	6	29	6	3	51
<i>Datomicra zosteræ</i> (Thomson,C.G.,1856)	2	0	0	0	0	2
<i>Atheta aquatica</i> (Thomson,C.G.,1852)	3	7	65	9	2	86
<i>Atheta aquatilis</i> (Thomson,C.G.,1867)	2	9	47	1	3	62
<i>Atheta brunneipennis</i> (Thomson,C.G.,1852)	2	5	18	5	1	31
<i>Atheta castanoptera</i> (Mannerheim,1830)	3	25	128	23	4	183
<i>Atheta graminicola</i> (Gravenhorst,1806)	33	29	199	46	10	317
<i>Atheta hypnorum</i> (Kiesenwetter,1850)	1	15	15	3	0	34
<i>Atheta incognita</i> (Sharp,1869)	0	5	2	0	0	7
<i>Atheta aeneicollis</i> (Sharp,1869)	35	35	99	24	4	197
<i>Atheta triangulum</i> (Kraatz,1856)	18	5	24	10	3	60
<i>Atheta xanthopus</i> (Thomson,C.G.,1856)	4	4	30	4	2	44
<i>Atheta basicornis</i> (Mulsant & Rey,1852)	0	0	1	2	1	4
<i>Dalotia coriaria</i> (Kraatz,1856)	7	6	48	10	0	71
<i>Atheta crassicornis</i> (Fabricius,1793)	35	54	238	50	18	395
<i>Atheta diversa</i> (Sharp,1869)	0	0	1	0	0	1
<i>Atheta fungicola</i> (Thomson,C.G.,1852)	0	4	6	1	0	11
<i>Atheta intermedia</i> (Thomson,C.G.,1852)	2	0	3	1	0	6
<i>Mycetota laticollis</i> (Stephens,1832)	41	13	199	38	4	295
<i>Atheta nidicola</i> (Johansen,1914)	0	1	2	2	0	5
<i>Atheta oblita</i> (Erichson,1839)	1	2	2	0	0	5
<i>Atheta paracrassicornis</i> Brundin,1954	0	0	0	1	0	1
<i>Atheta pilicornis</i> (Thomson,C.G.,1852)	0	1	0	1	0	2
<i>Atheta ravilla</i> (Erichson,1839)	12	8	42	8	2	72
<i>Atheta britanniae</i> (Bernhauer & Scheerpeltz,1926)	3	6	19	4	2	34
<i>Atheta strandiella</i> (Brundin,1954)	2	0	0	0	0	2
<i>Dimetrota atramentaria</i> (Gyllenhal,1810)	23	37	168	27	19	274
<i>Dimetrota cauta</i> (Erichson,1837)	4	0	1	1	0	6
<i>Dimetrota cinnamoptera</i> (Thomson,C.G.,1856)	0	12	29	3	6	50
<i>Dimetrota ischnocera</i> (Thomson,C.G.,1870)	2	6	26	3	1	38
<i>Dimetrota laevana</i> (Mulsant & Rey,1852)	0	5	10	5	2	22
<i>Badura macrocera</i> (Thomson,C.G.,1856)	4	2	15	6	2	29
<i>Dimetrota marcida</i> (Erichson,1837)	0	7	79	4	1	91
<i>Dimetrota nigripes</i> (Thomson,C.G.,1856)	13	7	48	2	6	76
<i>Dimetrota aeneipennis</i> (Thomson,C.G.,1856)	0	0	3	0	0	3
<i>Dimetrota setigera</i> (Sharp,1869)	4	3	28	2	1	38

Yorkshire Coleoptera checklist 2011

<i>Chaetida longicornis</i> (Gravenhorst,1802)	31	24	142	31	9	237
<i>Brundinia marina</i> (Mulsant & Rey,1853)	19	0	0	0	0	19
<i>Thinobaena vestita</i> (Gravenhorst,1806)	34	13	0	0	0	47
<i>Alevonota rufotestacea</i> (Kraatz,1856)	0	1	0	1	0	2
<i>Halobrecta algae</i> (Hardy,1851)	3	0	0	0	0	3
<i>Halobrecta flavipes</i> Thomson,C.G.,1861	6	2	0	0	0	8
<i>Alianta incana</i> (Erichson,1837)	8	1	32	5	2	48
<i>Pachnida nigella</i> (Erichson,1837)	21	1	53	6	1	82
<i>Thamiaraea cinnamomea</i> (Gravenhorst,1802)	0	0	1	0	0	1
<i>Thamiaraea hospita</i> (Markel,1844)	1	0	0	0	0	1
<i>Drusilla canaliculata</i> (Fabricius,1787)	33	43	67	10	5	158
<i>Pella humeralis</i> (Gravenhorst,1802)	2	3	5	0	1	11
<i>Lomechusa emarginata</i> (Paykull,1789)	0	0	6	0	0	6
<i>Dinarda dentata</i> (Gravenhorst,1806)	0	1	0	0	0	1
<i>Dinarda maerkeli</i> Kiesenwetter,1843	1	7	1	0	0	9
<i>Trichiusa immigrata</i> Lohse,1984	8	0	35	0	1	44
<i>Phloeopora corticalis</i> (Gravenhorst,1802)	1	2	7	2	1	13
<i>Phloeopora testacea</i> (Mannerheim,1830)	3	10	37	7	0	57
<i>Ilyobates nigricollis</i> (Paykull,1800)	1	0	1	3	0	5
<i>Ilyobates propinquus</i> (Aube,1850)	1	0	1	0	0	2
<i>Ilyobates bennetti</i> Donisthorpe,1914	4	0	2	2	0	8
<i>Calodera aethiops</i> (Gravenhorst,1802)	23	5	16	6	1	51
<i>Calodera nigrita</i> Mannerheim,1830	2	1	1	1	1	6
<i>Calodera protensa</i> Mannerheim,1830	0	0	0	2	0	2
<i>Calodera riparia</i> Erichson,1837	4	0	10	2	0	16
<i>Tetralaucopora longitarsis</i> (Erichson,1837)	24	17	25	11	3	80
<i>Tetralaucopora rubicunda</i> (Erichson,1837)	1	1	0	0	1	3
<i>Amarochara umbrosa</i> (Erichson,1837)	0	2	1	0	0	3
<i>Ocalea badia</i> Erichson,1837	1	3	1	1	0	6
<i>Ocalea latipennis</i> Sharp,1870	1	1	2	0	0	4
<i>Ocalea picata</i> (Stephens,1832)	4	19	45	11	6	85
<i>Meotica anglica</i> Benick in Muona,1991	1	0	0	0	0	1
<i>Meotica filiformis</i> (Motschulsky,1860)	4	0	10	2	0	16
<i>Meotica exilis</i> (Knoch in Gravenhorst,1806)	4	0	0	1	1	6
<i>Meotica exillima</i> Sharp,1915	1	1	3	1	0	6
<i>Ocyusa picina</i> (Aube,1850)	1	1	7	1	0	10
<i>Cousya longitarsis</i> (Thomson,C.G.,1867)	0	11	0	0	0	11
<i>Ocyusa maura</i> (Erichson,1837)	9	0	25	5	0	39
<i>Cousya nitidiventris</i> (Fagel,1958)	0	0	1	0	0	1
<i>Mniusa incrassata</i> (Mulsant & Rey,1852)	6	44	29	10	4	93
<i>Oxypoda alternans</i> (Gravenhorst,1802)	8	27	125	26	3	189
<i>Oxypoda flavicornis</i> Kraatz,1856	3	0	4	1	0	8
<i>Oxypoda annularis</i> Mannerheim,1830	6	25	22	6	0	59
<i>Oxypoda brachyptera</i> (Stephens,1832)	15	0	4	2	1	22
<i>Oxypoda elongatula</i> Aube,1850	30	23	241	35	8	337
<i>Oxypoda exoleta</i> Erichson,1839	8	1	3	0	0	12
<i>Oxypoda formiceticola</i> Markel,1841	1	5	9	0	0	15
<i>Oxypoda haemorrhoea</i> (Mannerheim,1830)	32	19	114	16	7	188
<i>Oxypoda induta</i> Mulsant & Rey,1861	1	2	1	0	0	4
<i>Oxypoda islandica</i> Kraatz,1857	0	0	0	2	0	2

Yorkshire Coleoptera checklist 2011

<i>Oxypoda lentula</i> Erichson,1837	20	1	5	0	0	26
<i>Oxypoda acuminata</i> (Stephens,1832)	3	7	22	3	3	38
<i>Oxypoda longipes</i> Mulsant & Rey,1861	2	0	1	0	0	3
<i>Oxypoda nigricornis</i> Motschulsky,1860	0	0	3	1	0	4
<i>Oxypoda opaca</i> (Gravenhorst,1802)	23	15	72	16	2	128
<i>Oxypoda procerula</i> Mannerheim,1830	4	18	28	2	1	53
<i>Oxypoda carbonaria</i> (Heer,1841)	2	5	19	1	0	27
<i>Oxypoda spectabilis</i> Markel,1844	1	6	2	2	0	11
<i>Oxypoda brevicornis</i> (Stephens,1832)	18	13	41	4	1	77
<i>Ischnoglossa prolixa</i> (Gravenhorst,1802)	1	0	11	0	0	12
<i>Dexiogyia corticina</i> (Erichson,1837)	0	1	0	0	0	1
<i>Thiasophila angulata</i> (Erichson,1837)	3	8	9	0	0	20
<i>Crataraea suturalis</i> (Mannerheim,1830)	1	1	5	2	1	10
<i>Haploglossa gentilis</i> (Markel,1844)	2	0	0	0	0	2
<i>Haploglossa nidicola</i> (Fairmaire,1852)	5	1	2	2	1	11
<i>Haploglossa villosula</i> (Stephens,1832)	2	7	1	3	1	14
<i>Tinotus morion</i> (Gravenhorst,1802)	10	13	21	9	2	55
<i>Aleochara bilineata</i> Gyllenhal,1810	7	4	5	0	0	16
<i>Aleochara binotata</i> Kraatz,1858	2	0	0	0	0	2
<i>Aleochara bipustulata</i> (Linnaeus,1761)	21	9	28	5	1	64
<i>Aleochara brevipennis</i> Gravenhorst,1806	0	0	1	0	0	1
<i>Aleochara cuniculorum</i> Kraatz,1858	5	0	1	2	1	9
<i>Aleochara curtula</i> (Goeze,1777)	22	15	46	18	11	112
<i>Aleochara discipennis</i> Mulsant & Rey,1853	0	0	2	0	0	2
<i>Aleochara kamila</i> Likovsky,1984	0	1	3	1	0	5
<i>Aleochara fumata</i> Gravenhorst,1802	0	0	2	0	0	2
<i>Aleochara funebris</i> Wollaston,1864	1	2	24	0	2	29
<i>Aleochara grisea</i> Kraatz,1856	12	3	0	0	0	15
<i>Aleochara inconspicua</i> Aube,1850	0	0	1	0	0	1
<i>Aleochara intricata</i> Mannerheim,1830	5	4	2	2	0	13
<i>Aleochara lanuginosa</i> Gravenhorst,1802	24	37	108	29	17	215
<i>Aleochara lygaea</i> Kraatz,1862	0	2	0	0	0	2
<i>Aleochara moerens</i> Gyllenhal,1827	0	0	2	0	0	2
<i>Aleochara moesta</i> Gravenhorst,1802	1	1	1	2	0	5
<i>Aleochara obscurella</i> Gravenhorst,1806	12	9	0	0	0	21
<i>Aleochara punctatella</i> Motschulsky,1858	5	3	0	0	0	8
<i>Aleochara ruficornis</i> Gravenhorst,1802	3	9	15	3	1	31
<i>Aleochara spadicea</i> (Erichson,1837)	8	3	1	5	0	17
<i>Aleochara sparsa</i> Heer,1839	8	10	7	2	3	30
<i>Aleochara verna</i> Say,1836	5	2	3	3	0	13
<i>Aleochara villosa</i> Mannerheim,1830	2	0	0	0	0	2
<i>Scaphidium quadrimaculatum</i> Olivier, 1790	1	16	35	2	0	54
<i>Scaphisoma agaricinum</i> (Linnaeus, 1758)	3	12	46	13	2	76
<i>Scaphisoma boleti</i> (Panzer, 1793)	8	12	10	6	0	36
<i>Siagonium quadricorne</i> Kirby, 1815	3	6	25	6	1	41
<i>Deleaster dichrous</i> (Gravenhorst, 1802)	1	6	3	5	5	20
<i>Syntomium aeneum</i> (Müller, P.W.J., 1821)	6	24	60	19	1	110
<i>Coprophilus striatulus</i> (Fabricius, 1792)	5	6	7	3	0	21
<i>Aploderus caelatus</i> (Gravenhorst, 1802)	1	4	4	1	1	11
<i>Bledius (Astycops) subterraneus</i> Erichson, 1839	0	14	3	7	2	26

Yorkshire Coleoptera checklist 2011

Bledius (Bledius) limicola Tottenham, 1940	6	0	1	0	0	7
Bledius (Bledius) spectabilis Kraatz, 1857	0	0	1	0	0	1
Bledius (Bledius) tricornis (Herbst, 1784)	3	1	0	0	0	4
Bledius (Dicarenus) fergussoni Joy, 1912	1	3	0	0	0	4
Bledius (Elbidus) bicornis (Germar, 1822)	0	1	0	0	0	1
Bledius (Hesperophilus) annae Sharp, 1911	2	4	2	11	1	20
Bledius (Hesperophilus) defensus Fauvel, 1872	0	2	1	9	2	14
Bledius (Hesperophilus) dissimilis Erichson, 1840	9	0	0	0	0	9
Bledius (Hesperophilus) erraticus Erichson, 1839	0	2	0	0	2	4
Bledius (Hesperophilus) femoralis (Gyllenhal, 1827)	0	3	0	0	0	3
Bledius (Hesperophilus) fuscipes Rye, 1865	0	0	0	1	0	1
Bledius (Hesperophilus) gallicus (Gravenhorst, 1806)	4	5	5	1	1	16
Bledius (Hesperophilus) longulus Erichson, 1839	2	4	0	1	2	9
Bledius (Hesperophilus) occidentalis Bondroit, 1907	1	0	1	0	0	2
Bledius (Hesperophilus) opacus (Block, 1799)	8	4	4	0	0	16
Bledius (Hesperophilus) pallipes (Gravenhorst, 1806)	0	0	0	1	0	1
Bledius (Hesperophilus) praetermissus Williams, 1929	6	0	1	0	0	7
Bledius (Hesperophilus) terebrans (Schiödte, 1866)	0	1	0	0	0	1
Carpelimus bilineatus Stephens, 1834	25	13	25	15	2	80
Carpelimus corticinus (Gravenhorst, 1806)	24	7	61	23	12	127
Carpelimus elongatulus (Erichson, 1839)	17	10	52	31	4	114
Carpelimus foveolatus (Sahlberg, C.R., 1832)	9	0	0	0	0	9
Carpelimus fuliginosus (Gravenhorst, 1802)	1	0	2	0	0	3
Carpelimus gracilis (Mannerheim, 1830)	0	0	2	0	0	2
Carpelimus halophilus (Kiesenwetter, 1844)	3	0	0	1	0	4
Carpelimus impressus (Boisduval & Lacordaire, 1835)	1	1	1	1	0	4
Carpelimus manchuricus subsp. subtilicornis (Roubal, 1946)	1	1	1	7	0	10
Carpelimus obesus (Kiesenwetter, 1844)	1	0	0	0	0	1
Carpelimus pusillus (Gravenhorst, 1802)	8	8	14	3	1	34
Carpelimus rivularis (Motschulsky, 1860)	25	8	48	18	7	106
Carpelimus zealandicus (Sharp, 1900)	6	2	10	3	0	21
Ochtheophilus andalusiacus (Fagel, 1957)	0	1	0	0	1	2
Ochtheophilus angustior (Bernhauer, 1943)	0	0	0	1	0	1
Ochtheophilus aureus (Fauvel, 1871)	0	3	2	2	1	8
Ochtheophilus omalinus (Erichson, 1840)	0	14	1	4	3	22
Thinobius bicolor Joy, 1911	0	2	0	1	0	3
Thinobius crinifer Smetana, 1959	0	1	0	1	0	2
Thinodromus arcuatus (Stephens, 1834)	1	7	1	4	2	15
Anotylus complanatus (Erichson, 1839)	14	20	44	8	4	90
Anotylus fairmairei (Pandellé, 1867)	0	1	2	0	0	3
Anotylus insecatus (Gravenhorst, 1806)	0	1	1	0	0	2
Anotylus inustus (Gravenhorst, 1806)	9	11	61	1	1	83
Anotylus maritimus Thomson, C.G., 1861	2	2	0	0	0	4
Anotylus mutator (Lohse, 1963)	2	4	21	5	0	32
Anotylus nitidulus (Gravenhorst, 1802)	1	8	7	4	1	21
Anotylus rugosus (Fabricius, 1775)	96	45	346	101	27	615
Anotylus saulcyi (Pandellé, 1867)	0	1	0	0	0	1
Anotylus sculpturatus (Gravenhorst, 1806)	38	35	218	69	16	376
Anotylus tetracarınatus (Block, 1799)	37	52	196	56	30	371
Oxytelus fulvipes Erichson, 1839	0	0	4	7	0	11

Yorkshire Coleoptera checklist 2011

<i>Oxytelus laqueatus</i> (Marsham, 1802)	15	25	111	38	14	203
<i>Oxytelus migrator</i> Fauvel, 1904	0	0	2	0	0	2
<i>Oxytelus piceus</i> (Linnaeus, 1767)	0	1	1	0	0	2
<i>Oxytelus sculptus</i> Gravenhorst, 1806	5	4	19	3	2	33
<i>Platystethus</i> (<i>Craetopycrus</i>) <i>cornutus</i> (Gravenhorst, 1802)	28	13	41	15	3	100
<i>Platystethus</i> (<i>Craetopycrus</i>) <i>nitens</i> (Sahlberg, C.R., 1832)	10	2	5	1	0	18
<i>Platystethus</i> (<i>Craetopycrus</i>) <i>nodifrons</i> (Mannerheim, 1830)	24	3	1	2	4	34
<i>Platystethus</i> (<i>Platystethus</i>) <i>arenarius</i> (Fourcroy, 1785)	23	28	98	17	12	178
<i>Oxyporus rufus</i> (Linnaeus, 1758)	13	6	16	4	0	39
<i>Stenus</i> (<i>Hemistenus</i>) <i>aceris</i> Stephens, 1833	1	5	6	1	2	15
<i>Stenus</i> (<i>Hemistenus</i>) <i>fuscicornis</i> Erichson, 1840	2	2	16	2	1	23
<i>Stenus</i> (<i>Hemistenus</i>) <i>geniculatus</i> Gravenhorst, 1806	0	0	1	0	0	1
<i>Stenus</i> (<i>Hemistenus</i>) <i>impressus</i> Germar, 1824	45	59	315	76	19	514
<i>Stenus</i> (<i>Hemistenus</i>) <i>ochropus</i> Kiesenwetter, 1858	1	0	2	1	0	4
<i>Stenus</i> (<i>Hemistenus</i>) <i>ossium</i> Stephens, 1833	10	6	16	5	0	37
<i>Stenus</i> (<i>Hemistenus</i>) <i>pallipes</i> Gravenhorst, 1802	9	0	8	1	0	18
<i>Stenus</i> (<i>Hemistenus</i>) <i>palustris</i> Erichson, 1839	9	2	6	5	0	22
<i>Stenus</i> (<i>Hemistenus</i>) <i>subaeneus</i> Erichson, 1840	6	3	11	1	0	21
<i>Stenus</i> (<i>Hypostenus</i>) <i>cicindeloides</i> (Schaller, 1783)	26	9	47	20	0	102
<i>Stenus</i> (<i>Hypostenus</i>) <i>fulvicornis</i> Stephens, 1833	25	15	114	30	4	188
<i>Stenus</i> (<i>Hypostenus</i>) <i>kiesenwetteri</i> Rosenhauer, 1856	0	0	1	0	0	1
<i>Stenus</i> (<i>Hypostenus</i>) <i>latifrons</i> Erichson, 1839	13	7	34	8	1	63
<i>Stenus</i> (<i>Hypostenus</i>) <i>oscillator</i> Rye, 1870	0	0	8	0	0	8
<i>Stenus</i> (<i>Hypostenus</i>) <i>similis</i> (Herbst, 1784)	61	58	201	49	6	375
<i>Stenus</i> (<i>Hypostenus</i>) <i>solutus</i> Erichson, 1840	1	1	17	2	0	21
<i>Stenus</i> (<i>Hypostenus</i>) <i>tarsalis</i> Ljungh, 1810	20	6	7	5	1	39
<i>Stenus</i> (<i>Metatesnus</i>) <i>bifoveolatus</i> Gyllenhal, 1827	13	6	47	22	1	89
<i>Stenus</i> (<i>Metatesnus</i>) <i>binotatus</i> Ljungh, 1804	2	10	22	1	0	35
<i>Stenus</i> (<i>Metatesnus</i>) <i>brevipennis</i> Thomson, C.G., 1851	1	8	6	4	5	24
<i>Stenus</i> (<i>Metatesnus</i>) <i>flavipes</i> Stephens, 1833	13	23	33	42	5	116
<i>Stenus</i> (<i>Metatesnus</i>) <i>nitidiusculus</i> Stephens, 1833	14	17	83	42	6	162
<i>Stenus</i> (<i>Metatesnus</i>) <i>niveus</i> Fauvel, 1865	2	0	1	0	0	3
<i>Stenus</i> (<i>Metatesnus</i>) <i>pallitarsis</i> Stephens, 1833	39	7	41	14	2	103
<i>Stenus</i> (<i>Metatesnus</i>) <i>picipennis</i> Erichson, 1840	5	4	25	6	1	41
<i>Stenus</i> (<i>Metatesnus</i>) <i>picipes</i> Stephens, 1833	17	12	54	28	6	117
<i>Stenus</i> (<i>Metatesnus</i>) <i>pubescens</i> Stephens, 1833	10	4	35	5	0	54
<i>Stenus</i> (<i>Metatesnus</i>) <i>umbratilis</i> Casey, 1884	0	0	0	5	1	6
<i>Stenus</i> (<i>Stenus</i>) <i>argus</i> Gravenhorst, 1806	10	2	0	3	0	15
<i>Stenus</i> (<i>Stenus</i>) <i>asphaltinus</i> Erichson, 1840	0	0	1	0	0	1
<i>Stenus</i> (<i>Stenus</i>) <i>ater</i> Mannerheim, 1830	3	0	4	2	0	9
<i>Stenus</i> (<i>Stenus</i>) <i>atratus</i> Erichson, 1839	4	5	9	5	0	23
<i>Stenus</i> (<i>Stenus</i>) <i>biguttatus</i> (Linnaeus, 1758)	14	11	13	9	5	52
<i>Stenus</i> (<i>Stenus</i>) <i>bimaculatus</i> Gyllenhal, 1810	37	19	77	57	8	198
<i>Stenus</i> (<i>Stenus</i>) <i>boops</i> Ljungh, 1810	38	13	79	29	3	162
<i>Stenus</i> (<i>Stenus</i>) <i>canaliculatus</i> Gyllenhal, 1827	14	1	28	15	1	59
<i>Stenus</i> (<i>Stenus</i>) <i>carbonarius</i> Gyllenhal, 1827	22	2	16	7	0	47
<i>Stenus</i> (<i>Stenus</i>) <i>clavicornis</i> (Scopoli, 1763)	67	34	240	44	10	395
<i>Stenus</i> (<i>Stenus</i>) <i>comma</i> LeConte, 1863	5	4	8	5	0	22
<i>Stenus</i> (<i>Stenus</i>) <i>europaeus</i> Puthz, 1966	22	3	13	9	1	48
<i>Stenus</i> (<i>Stenus</i>) <i>fuscipes</i> Gravenhorst, 1802	0	0	1	1	0	2

Yorkshire Coleoptera checklist 2011

Stenus (Stenus) guttula Müller, P.W.J., 1821	15	14	9	10	7	55
Stenus (Stenus) guynemeri Jacquelin du Val, 1850	0	5	20	7	9	41
Stenus (Stenus) incrassatus Erichson, 1839	1	0	1	0	0	2
Stenus (Stenus) junco (Paykull, 1789)	79	27	186	71	10	373
Stenus (Stenus) lustrator Erichson, 1839	2	2	6	0	0	10
Stenus (Stenus) melanarius Stephens, 1833	4	0	12	4	0	20
Stenus (Stenus) melanopus (Marsham, 1802)	4	0	2	0	0	6
Stenus (Stenus) morio Gravenhorst, 1806	1	1	0	0	0	2
Stenus (Stenus) nanus Stephens, 1833	6	11	54	9	4	84
Stenus (Stenus) nitens Stephens, 1833	3	1	23	1	0	28
Stenus (Stenus) providus Erichson, 1839	25	25	93	36	9	188
Stenus (Stenus) pusillus Stephens, 1833	21	10	49	25	5	110
Stenus (Tesnus) brunnipes Stephens, 1833	67	22	191	63	14	357
Stenus (Tesnus) crassus Stephens, 1833	3	1	8	5	0	17
Stenus (Tesnus) formicetorum Mannerheim, 1843	2	2	1	0	0	5
Stenus (Tesnus) nigritulus Gyllenhal, 1827	9	3	7	1	1	21
Dianous coerulescens (Gyllenhal, 1810)	1	17	21	18	18	75
Euaesthetus bipunctatus (Ljungh, 1804)	2	1	7	14	2	26
Euaesthetus laeviusculus Mannerheim, 1844	2	4	17	4	0	27
Euaesthetus ruficapillus Lacordaire, 1835	16	4	26	6	3	55
Pseudopsis sulcata Newman, 1834	1	3	4	3	0	11
Achenium depressum (Gravenhorst, 1802)	1	0	0	0	0	1
Achenium humile (Nicolai, 1822)	1	0	0	0	0	1
Astenus (Astenognathus) procerus (Gravenhorst, 1806)	0	0	1	0	0	1
Astenus (Astenognathus) pulchellus (Heer, 1839)	6	1	10	2	0	19
Astenus (Astenus) lyonessius (Joy, 1908)	14	1	5	3	0	23
Lathrobium (Tetartoepus) quadratum (Paykull, 1789)	16	3	24	5	0	48
Lathrobium (Tetartoepus) terminatum Gravenhorst, 1802	6	6	71	8	3	94
Lathrobium (Tetartoepus) zetterstedti Rye, 1872	0	1	3	1	2	7
Lathrobium (Lathrobium) brunnipes (Fabricius, 1792)	49	36	272	92	25	474
Lathrobium (Lathrobium) elongatum (Linnaeus, 1767)	9	4	13	9	3	38
Lathrobium (Lathrobium) fovulum Stephens, 1833	1	0	2	5	0	8
Lathrobium (Lathrobium) fulvipenne (Gravenhorst, 1806)	25	21	75	19	3	143
Lathrobium (Lathrobium) geminum Kraatz, 1857	13	8	58	9	5	93
Lathrobium (Lathrobium) impressum Heer, 1841	7	1	11	13	0	32
Lathrobium (Lathrobium) longulum Gravenhorst, 1802	7	2	10	1	1	21
Lathrobium (Lathrobium) pallidipenne Hichhuth, 1851	1	4	1	1	0	7
Lathrobium (Lathrobium) rufipenne Gyllenhal, 1813	0	0	1	0	0	1
Lithocharis nigriceps Kraatz, 1859	15	2	40	7	1	65
Lithocharis ochracea (Gravenhorst, 1802)	11	2	20	5	2	40
Lobrathium multipunctum (Gravenhorst, 1802)	1	6	5	1	1	14
Medon apicalis (Kraatz, 1857)	1	0	0	0	0	1
Medon brunneus (Erichson, 1839)	0	1	2	0	0	3
Medon castaneus (Gravenhorst, 1802)	0	0	1	0	0	1
Medon fuscus (Mannerheim, 1830)	0	4	0	0	0	4
Medon pocofer (Peyron, 1857)	0	1	0	0	0	1
Ochtheophilum fracticorne (Paykull, 1800)	4	7	11	6	1	29
Paederus fuscipes Curtis, 1826	0	0	1	0	0	1
Paederus littoralis Gravenhorst, 1802	1	3	1	1	0	6
Paederus riparius (Linnaeus, 1758)	3	8	3	1	0	15

Yorkshire Coleoptera checklist 2011

<i>Pseudomedon obscurellus</i> (Erichson, 1840)	0	0	2	1	0	3
<i>Pseudomedon obsoletus</i> (Nordmann, 1837)	0	0	7	0	0	7
<i>Rugilus erichsoni</i> (Fauvel, 1867)	13	5	28	17	0	63
<i>Rugilus orbiculatus</i> (Paykull, 1789)	46	19	158	29	2	254
<i>Rugilus rufipes</i> Germar, 1835	44	7	82	28	4	165
<i>Rugilus subtilis</i> (Erichson, 1840)	0	1	0	0	0	1
<i>Scopaeus sulcicollis</i> (Stephens, 1833)	0	0	1	0	0	1
<i>Sunius bicolor</i> (Olivier, 1795)	4	0	1	0	0	5
<i>Sunius melanocephalus</i> (Fabricius, 1792)	3	1	0	1	0	5
<i>Sunius propinquus</i> (Brisout, 1867)	15	2	16	0	0	33
<i>Bisnius cephalotes</i> (Gravenhorst, 1802)	12	8	57	14	3	94
<i>Bisnius fimetarius</i> (Gravenhorst, 1802)	37	22	272	36	8	375
<i>Bisnius nigriventris</i> (Thomson, C.G., 1867)	4	1	8	3	2	18
<i>Bisnius parvus</i> (Sharp, 1874)	4	0	13	2	0	19
<i>Bisnius puella</i> (Nordmann, 1837)	2	3	29	10	2	46
<i>Bisnius sordidus</i> (Gravenhorst, 1802)	22	5	67	11	0	105
<i>Bisnius subuliformis</i> (Gravenhorst, 1802)	1	0	3	2	0	6
<i>Cafius fucicola</i> Curtis, 1830	1	1	0	0	0	2
<i>Cafius xantholoma</i> (Gravenhorst, 1806)	13	9	2	0	0	24
<i>Erichsonius cinerascens</i> (Gravenhorst, 1802)	2	4	10	0	0	16
<i>Gabrius appendiculatus</i> Sharp, 1910	1	3	8	4	1	17
<i>Gabrius astutoides</i> (Strand, A., 1946)	1	0	0	0	0	1
<i>Gabrius bishopi</i> Sharp, 1910	12	0	4	4	1	21
<i>Gabrius breviventer</i> (Sperk, 1835)	22	17	93	35	3	170
<i>Gabrius nigritulus</i> (Gravenhorst, 1802)	6	7	14	4	1	32
<i>Gabrius osseticus</i> (Kolenati, 1846)	0	1	0	0	0	1
<i>Gabrius piliger</i> Mulsant & Rey, 1876	5	0	36	5	2	48
<i>Gabrius splendidulus</i> (Gravenhorst, 1802)	6	13	80	14	6	119
<i>Gabrius trossulus</i> (Nordmann, 1837)	17	6	29	26	1	79
<i>Gabrius velox</i> Sharp, 1910	0	1	0	0	0	1
<i>Gabronthus thermarum</i> (Aubé, 1850)	1	0	0	0	0	1
<i>Neobisnius lathrobioides</i> (Baudi, 1848)	0	0	1	0	0	1
<i>Neobisnius procerulus</i> (Gravenhorst, 1806)	1	0	2	0	0	3
<i>Neobisnius prolixus</i> (Erichson, 1840)	0	0	1	0	0	1
<i>Neobisnius villosulus</i> (Stephens, 1833)	0	1	1	0	0	2
<i>Philonthus addendus</i> Sharp, 1867	2	5	10	2	2	21
<i>Philonthus albipes</i> (Gravenhorst, 1802)	8	3	22	3	1	37
<i>Philonthus atratus</i> (Gravenhorst, 1802)	3	4	5	0	0	12
<i>Philonthus carbonarius</i> (Gravenhorst, 1802)	31	47	160	39	7	284
<i>Philonthus cognatus</i> Stephens, 1832	33	34	132	55	8	262
<i>Philonthus concinnus</i> (Gravenhorst, 1802)	12	1	21	5	0	39
<i>Philonthus corruscus</i> (Gravenhorst, 1802)	0	0	0	1	0	1
<i>Philonthus corvinus</i> Erichson, 1839	3	3	0	0	0	6
<i>Philonthus cruentatus</i> (Gmelin in Linnaeus, 1790)	4	2	7	1	0	14
<i>Philonthus debilis</i> (Gravenhorst, 1802)	4	2	30	3	0	39
<i>Philonthus decorus</i> (Gravenhorst, 1802)	7	22	176	34	15	254
<i>Philonthus discoideus</i> (Gravenhorst, 1802)	4	5	10	2	0	21
<i>Philonthus ebeninus</i> (Gravenhorst, 1802)	0	4	5	2	0	11
<i>Philonthus fumarius</i> (Gravenhorst, 1806)	0	1	0	6	0	7
<i>Philonthus intermedius</i> (Lacordaire, 1835)	12	10	38	5	2	67

Yorkshire Coleoptera checklist 2011

Philonthus jurgans Tottenham, 1937	2	2	6	2	0	12
Philonthus laminatus (Creutzer, 1799)	17	16	84	27	8	152
Philonthus longicornis Stephens, 1832	11	5	10	5	1	32
Philonthus mannerheimi Fauvel, 1869	0	2	0	0	0	2
Philonthus marginatus (Müller, O.F., 1764)	16	14	73	27	6	136
Philonthus micans (Gravenhorst, 1802)	19	14	7	6	2	48
Philonthus micantoides Benick & Lohse, 1956	11	2	12	8	0	33
Philonthus nigrita (Gravenhorst, 1806)	2	9	4	4	0	19
Philonthus parvicornis (Gravenhorst, 1802)	0	0	3	2	0	5
Philonthus politus (Linnaeus, 1758)	14	5	43	9	3	74
Philonthus quisquiliarius (Gyllenhal, 1810)	26	9	34	14	6	89
Philonthus rectangulus Sharp, 1874	4	5	23	7	2	41
Philonthus rotundicollis (Ménétriés, 1832)	2	10	23	5	0	40
Philonthus rubripennis Stephens, 1832	0	20	0	2	0	22
Philonthus rufipes (Stephens, 1832)	0	0	1	0	0	1
Philonthus sanguinolentus (Gravenhorst, 1802)	9	13	26	2	0	50
Philonthus spinipes Sharp, 1874	0	0	3	0	0	3
Philonthus splendens (Fabricius, 1792)	16	10	37	11	11	85
Philonthus succicola Thomson, C.G., 1860	17	9	90	19	8	143
Philonthus tenuicornis Mulsant & Rey, 1853	11	7	74	11	2	105
Philonthus umbratilis (Gravenhorst, 1802)	8	3	28	2	0	41
Philonthus varians (Paykull, 1789)	50	35	158	49	19	311
Philonthus ventralis (Gravenhorst, 1802)	1	3	6	3	0	13
Remus sericeus Holme, 1837	0	1	0	0	0	1
Creophilus maxillosus (Linnaeus, 1758)	13	8	29	4	6	60
Dinothenarus pubescens (De Geer, 1774)	3	1	5	3	0	12
Ocypus (Matidus) brunnipes (Fabricius, 1781)	25	13	50	14	3	105
Ocypus (Matidus) nitens (Schrank, 1781)	1	1	4	0	0	6
Ocypus (Ocypus) olens (Müller, O.F., 1764)	33	8	132	28	1	202
Ocypus (Ocypus) ophthalmicus (Scopoli, 1763)	0	0	8	0	0	8
Ocypus (Pseudocypus) aeneocephalus (De Geer, 1774)	5	13	40	32	7	97
Ocypus (Pseudocypus) fortunatarum (Wollaston, 1871)	0	1	1	0	0	2
Ocypus (Pseudocypus) fuscatus (Gravenhorst, 1802)	2	2	7	1	0	12
Ontholestes murinus (Linnaeus, 1758)	3	0	14	1	0	18
Ontholestes tessellatus (Geoffroy, 1785)	1	2	20	7	1	31
Platydracus fulvipes (Scopoli, 1763)	1	0	1	0	1	3
Platydracus latebricola (Gravenhorst, 1806)	0	4	0	0	0	4
Platydracus stercorarius (Olivier, 1795)	8	6	20	5	6	45
Staphylinus caesareus Cederhjelm, 1798	0	0	1	0	0	1
Staphylinus dimidiaticornis Gemminger, 1851	0	1	0	0	1	2
Staphylinus erythropterus Linnaeus, 1758	2	11	0	1	3	17
Tasgius (Rayacheila) globulifer (Geoffroy, 1785)	7	7	21	6	2	43
Tasgius (Rayacheila) melanarius (Heer, 1839)	9	4	46	20	2	81
Tasgius (Rayacheila) morsitans (Rossi, 1790)	8	4	11	1	2	26
Tasgius (Rayacheila) winkleri Bernhauer, 1906	3	3	3	0	0	9
Tasgius (Tasgius) ater (Gravenhorst, 1802)	24	2	22	2	0	50
Tasgius (Tasgius) pedator (Gravenhorst, 1802)	0	1	0	1	0	2
Heterothops binotatus (Gravenhorst, 1802)	5	2	0	2	0	9
Heterothops dissimilis (Gravenhorst, 1802)	6	3	11	1	0	21
Heterothops minutus Wollaston, 1860	3	0	10	1	0	14

Yorkshire Coleoptera checklist 2011

Heterothops niger Kraatz, 1868	3	1	2	1	0	7
Heterothops praeius Erichson, 1839	1	1	1	2	0	5
Quedius (Distichalius) cinctus (Paykull, 1790)	9	3	20	10	0	42
Quedius (Microsaurus) brevicornis (Thomson, C.G., 1860)	4	0	0	1	0	5
Quedius (Microsaurus) brevis Erichson, 1840	0	4	2	0	1	7
Quedius (Microsaurus) cruentus (Olivier, 1795)	10	12	59	15	3	99
Quedius (Microsaurus) fulgidus (Fabricius, 1792)	1	2	1	1	0	5
Quedius (Microsaurus) invreae Gridelli, 1924	0	2	9	3	0	14
Quedius (Microsaurus) lateralis (Gravenhorst, 1802)	1	7	30	4	1	43
Quedius (Microsaurus) longicornis Kraatz, 1857	3	3	4	1	0	11
Quedius (Microsaurus) maurus (Sahlberg, C.R., 1830)	3	9	17	3	0	32
Quedius (Microsaurus) mesomelinus (Marsham, 1802)	16	11	89	14	5	135
Quedius (Microsaurus) nigrocaeruleus Fauvel, 1874	3	0	0	2	0	5
Quedius (Microsaurus) puncticollis (Thomson, C.G., 1867)	1	4	1	6	1	13
Quedius (Microsaurus) scitus (Gravenhorst, 1806)	7	6	24	7	2	46
Quedius (Microsaurus) truncicola Fairmaire & Laboulbène, 1856	0	2	4	1	0	7
Quedius (Microsaurus) xanthopus Erichson, 1839	0	9	3	0	1	13
Quedius (Quedionuchus) plagiatus Mannerheim, 1843	0	5	8	11	1	25
Quedius (Quedius) curtipennis Bernhauer, 1908	19	23	165	58	11	276
Quedius (Quedius) fuliginosus (Gravenhorst, 1802)	14	9	56	21	4	104
Quedius (Quedius) levicollis (Brullé, 1832)	12	6	47	13	4	82
Quedius (Quedius) molochinus (Gravenhorst, 1806)	15	24	83	25	12	159
Quedius (Quedius) simplicifrons Fairmaire, 1861	16	0	0	0	0	16
Quedius (Raphirus) auricomus Kiesenwetter, 1850	0	5	6	6	5	22
Quedius (Raphirus) boopoides Munster, 1923	4	0	4	1	1	10
Quedius (Raphirus) boops (Gravenhorst, 1802)	14	19	43	13	7	96
Quedius (Raphirus) fulvicollis (Stephens, 1833)	0	5	5	4	1	15
Quedius (Raphirus) fumatus (Stephens, 1833)	2	11	41	12	1	67
Quedius (Raphirus) humeralis Stephens, 1832	4	0	4	1	0	9
Quedius (Raphirus) maurorufus (Gravenhorst, 1806)	29	7	127	44	7	214
Quedius (Raphirus) nemoralis Baudi, 1848	4	0	21	1	0	26
Quedius (Raphirus) nigriceps Kraatz, 1857	1	14	29	3	0	47
Quedius (Raphirus) nitipennis (Stephens, 1833)	8	8	14	13	6	49
Quedius (Raphirus) persimilis Mulsant & Rey, 1876	2	0	23	3	1	29
Quedius (Raphirus) picipes (Mannerheim, 1830)	6	9	16	6	1	38
Quedius (Raphirus) riparius Kellner, 1843	1	1	1	0	0	3
Quedius (Raphirus) schatzmayri Gridelli, 1922	4	5	19	6	0	34
Quedius (Raphirus) scintillans (Gravenhorst, 1806)	11	2	58	22	3	96
Quedius (Raphirus) semiaeneus (Stephens, 1833)	8	5	7	2	1	23
Quedius (Raphirus) semiobscurus (Marsham, 1802)	15	6	25	8	0	54
Quedius (Raphirus) umbrinus Erichson, 1839	5	12	12	13	5	47
Atrecus affinis (Paykull, 1789)	5	19	92	18	5	139
Othius angustus Stephens, 1833	4	5	33	7	7	56
Othius laeviusculus Stephens, 1833	10	3	18	1	2	34
Othius punctulatus (Goeze, 1777)	13	14	118	41	9	195
Othius subuliformis Stephens, 1833	17	29	101	19	12	178
Gauropterus fulgidus (Fabricius, 1787)	2	1	3	3	0	9
Gyrohypnus angustatus Stephens, 1833	19	13	39	18	1	90
Gyrohypnus atratus (Heer, 1839)	0	3	0	0	0	3
Gyrohypnus fracticornis (Müller, O.F., 1776)	23	11	91	19	4	148

Yorkshire Coleoptera checklist 2011

Gyrophypnus punctulatus (Paykull, 1789)	1	1	47	8	1	58
Leptacinus batychrus (Gyllenhal, 1827)	0	4	14	0	0	18
Leptacinus formicetorum Märkel, 1841	0	5	7	0	0	12
Leptacinus intermedius Donisthorpe, 1936	1	0	1	2	0	4
Leptacinus pusillus (Stephens, 1833)	11	3	64	10	1	89
Megalinus glabratus (Gravenhorst, 1802)	34	12	25	10	0	81
Nudobius lentus (Gravenhorst, 1806)	2	9	12	7	1	31
Phacophallus parumpunctatus (Gyllenhal, 1827)	6	2	8	1	0	17
Xantholinus (Acanthophallus) laevigatus Jacobsen, 1849	0	0	1	1	0	2
Xantholinus (Purrolinus) elegans (Olivier, 1795)	8	1	17	1	1	28
Xantholinus (Purrolinus) tricolor (Fabricius, 1787)	1	0	0	0	0	1
Xantholinus (Xantholinus) gallicus Coiffait, 1956	0	3	0	0	1	4
Xantholinus (Xantholinus) linearis (Olivier, 1795)	85	41	281	56	17	480
Xantholinus (Xantholinus) longiventris Heer, 1839	20	20	66	27	0	133

Lucanidae [4/3]

Sinodendron cylindricum (Linnaeus, 1758)	8	24	98	25	5	160
Lucanus cervus (Linnaeus, 1758)	1	2	3	3	0	9
Dorcus parallelipedus (Linnaeus, 1758)	22	3	36	7	0	68

Trogidae [3/2]

Trox sabulosus (Linnaeus, 1758)	0	0	0	1	0	1
Trox scaber (Linnaeus, 1767)	0	1	10	2	0	13

Bolboceratidae [1/1]

Odonteus armiger (Scopoli, 1772)	0	0	2	0	0	2
----------------------------------	---	---	---	---	---	---

Geotrupidae [7/6]

Typhaeus typhoeus (Linnaeus, 1758)	13	11	3	1	1	29
Anoplotrupes stercorosus (Hartmann in Scriba, 1791)	0	5	7	0	1	13
Geotrupes mutator (Marsham, 1802)	2	0	0	0	0	2
Geotrupes spiniger (Marsham, 1802)	10	1	12	3	1	27
Geotrupes stercorarius (Linnaeus, 1758)	6	6	40	6	8	66
Trypocopris vernalis (Linnaeus, 1758)	1	4	0	0	1	6

Scarabiidae [83/57]

Onthophagus (Paleonthophagus) fracticornis (Preyssler, 1790)	0	0	1	0	0	1
Onthophagus (Paleonthophagus) joannae Goljan, 1953	0	0	0	1	0	1
Onthophagus (Paleonthophagus) nuchicornis (Linnaeus, 1758)	0	1	0	0	0	1
Onthophagus (Paleonthophagus) similis (Scriba, 1790)	0	0	3	0	0	3
Onthophagus (Paleonthophagus) vacca (Linnaeus, 1767)	0	0	1	0	0	1
Aegialia (Aegialia) arenaria (Fabricius, 1787)	12	5	0	0	0	17
Aegialia (Psammoporus) insularis Pittino, 2006	1	6	1	3	3	14
Euheptaulacus villosus (Gyllenhal in Schönherr, 1806)	0	1	1	2	0	4
Heptaulacus testudinarius (Fabricius, 1775)	0	2	0	2	0	4
Oxyomus sylvestris (Scopoli, 1763)	0	2	3	0	0	5
Aphodius (Acrossus) depressus (Kugelann, 1792)	8	11	30	11	4	64
Aphodius (Acrossus) luridus (Fabricius, 1775)	4	5	12	2	0	23
Aphodius (Acrossus) rufipes (Linnaeus, 1758)	20	15	92	41	161	329
Aphodius (Agoliinus) lapponum Gyllenhal, 1808	0	6	23	6	2	37

Yorkshire Coleoptera checklist 2011

Aphodius (Agrilinus) ater (De Geer, 1774)	19	21	61	19	10	130
Aphodius (Agrilinus) constans Duftschmid, 1805	0	5	3	5	3	16
Aphodius (Agrilinus) rufus (Moll, 1782)	11	6	25	3	11	56
Aphodius (Agrilinus) sordidus (Fabricius, 1775)	1	1	4	0	0	6
Aphodius (Ammoecius) brevis (Erichson, 1848)	0	0	0	1	0	1
Aphodius (Aphodius) fimetarius (Linnaeus, 1758)	7	17	43	5	5	77
Aphodius (Aphodius) foetens (Fabricius, 1787)	4	6	11	5	0	26
Aphodius (Aphodius) foetidus (Herbst, 1783)	3	3	0	0	1	7
Aphodius (Bodilus) ictericus (Laicharting, 1781)	2	0	0	1	0	3
Aphodius (Calomosternus) granarius (Linnaeus, 1767)	9	6	10	2	1	28
Aphodius (Chilothorax) conspurcatus (Linnaeus, 1758)	0	3	5	6	0	14
Aphodius (Chilothorax) distinctus (Müller, O.F., 1776)	5	4	8	2	0	19
Aphodius (Chilothorax) paykulli Bedel, 1907	1	0	7	0	0	8
Aphodius (Colobopterus) erraticus (Linnaeus, 1758)	2	3	0	1	0	6
Aphodius (Esymus) merdarius (Fabricius, 1775)	3	4	3	1	0	11
Aphodius (Esymus) pusillus (Herbst, 1789)	2	2	8	4	2	18
Aphodius (Euorodalus) coenosus (Panzer, 1798)	0	0	1	0	0	1
Aphodius (Eupleurus) subterraneus (Linnaeus, 1758)	1	1	4	2	0	8
Aphodius (Limarus) zenkeri Germar, 1813	1	0	3	0	0	4
Aphodius (Liothorax) plagiatus (Linnaeus, 1767)	4	0	0	0	0	4
Aphodius (Melinopterus) prodromus (Brahm, 1790)	18	18	48	12	5	101
Aphodius (Melinopterus) sphacelatus (Panzer, 1798)	7	16	68	8	6	105
Aphodius (Nimbus) contaminatus (Herbst, 1783)	9	12	46	6	4	77
Aphodius (Nimbus) obliteratus Panzer, 1823	1	2	2	5	0	10
Aphodius (Otophorus) haemorrhoidalis (Linnaeus, 1758)	6	5	9	2	1	23
Aphodius (Plagiogonus) arenarius (Olivier, 1789)	1	0	4	2	1	8
Aphodius (Planolinus) borealis Gyllenhal, 1827	0	3	5	2	2	12
Aphodius (Planolinus) fasciatus (Olivier, 1789)	0	7	2	1	5	15
Aphodius (Sigorus) porcus (Fabricius, 1792)	0	1	0	1	0	2
Aphodius (Teuchestes) fossor (Linnaeus, 1758)	19	20	40	8	9	96
Aphodius (Volinus) sticticus (Panzer, 1798)	1	0	1	2	1	5
Psammodius asper (Fabricius, 1775)	0	1	0	0	0	1
Melolontha hippocastani Fabricius, 1801	0	0	1	0	0	1
Melolontha melolontha (Linnaeus, 1758)	5	9	41	23	65	143
Polyphylla fullo (Linnaeus, 1758)	0	0	1	0	0	1
Amphimallon solstitiale (Linnaeus, 1758)	1	4	1	0	0	6
Serica brunnea (Linnaeus, 1758)	9	6	39	20	20	94
Hoplia philanthus (Füessly, 1775)	0	5	3	10	1	19
Anomala dubia (Scopoli, 1763)	15	3	1	0	0	19
Phyllopertha horticola (Linnaeus, 1758)	12	4	9	1	0	26
Oxythyrea funesta (Poda, 1761)	0	0	2	0	0	2
Cetonia aurata (Linnaeus, 1758)	1	3	1	0	0	5
Protaetia (Potosia) metallica (Herbst, 1782)	0	1	0	0	0	1
<u>Clambidae [10/10]</u>						
Calyptomerus dubius (Marsham, 1802)	18	3	16	3	0	40
Clambus Fischer von Waldheim, 1823/4	0	0	1	0	0	1
Clambus armadillo (De Geer, 1774)	7	8	4	3	2	24
Clambus evae Endrödy-Younga, 1960	0	1	0	0	1	2
Clambus gibbulus (LeConte, 1850)	0	0	4	0	0	4

Yorkshire Coleoptera checklist 2011

Clambus nigrellus Reitter, 1914	0	0	0	4	1	5
Clambus nigriclavus Stephens, 1835	0	0	0	4	0	4
Clambus pubescens Redtenbacher, 1849	12	4	17	1	1	35
Clambus punctulum (Beck, 1817)	3	3	17	1	0	24
Clambus simsoni Blackburn, 1902	0	0	4	0	0	4

Scirtidae [20/18]

Elodes elongata Tournier, 1868	0	0	2	2	1	5
Elodes minuta (Linnaeus, 1767)	4	11	12	9	3	39
Elodes pseudominuta Klausnitzer, 1971	1	12	31	1	4	49
Elodes tricuspidata Nyholm, 1985	0	0	0	1	0	1
Odeles marginata (Fabricius, 1798)	2	13	41	7	10	73
Microcara testacea (Linnaeus, 1767)	40	9	68	29	2	148
Cyphon coarctatus Paykull, 1799	44	53	172	65	13	347
Cyphon hilaris Nyholm, 1844	4	7	30	4	1	46
Cyphon laevipennis Tournier, 1868	25	3	59	8	1	96
Cyphon ochraceus Stephens, 1830	16	13	27	12	1	69
Cyphon padi (Linnaeus, 1758)	18	16	81	27	6	148
Cyphon palustris Thomson, C.G., 1855	2	6	22	4	5	39
Cyphon pubescens (Fabricius, 1792)	1	0	1	1	0	3
Cyphon punctipennis Sharp, 1873	1	2	0	0	0	3
Cyphon variabilis (Thunberg, 1787)	14	12	87	18	2	133
Prionocyphon serricornis (Müller, P.W.J., 1821)	0	5	6	1	0	12
Hydrocyphon deflexicollis (Müller, P.W.J., 1821)	0	4	0	0	3	7
Scirtes hemisphaericus (Linnaeus, 1758)	12	3	18	12	1	46

Dascillidae [1/1]

Dascillus cervinus (Linnaeus, 1758)	5	12	11	14	6	48
-------------------------------------	---	----	----	----	---	----

Buprestidae [14/6]

Agrilus (Agrilus) viridis (Linnaeus, 1758)	0	2	0	0	0	2
Agrilus (Anambus) angustulus (Illiger, 1803)	3	2	1	0	0	6
Agrilus (Anambus) laticornis (Illiger, 1803)	1	7	7	2	0	17
Aphanisticus pusillus (Olivier, 1790)	0	1	0	0	0	1
Trachys minuta (Linnaeus, 1758)	2	0	0	0	0	2
Trachys troglodytes Gyllenhal in Schoenherr, 1817	1	1	0	0	0	2

Byrrhidae [13/11]

Simplocaria maculosa Erichson, 1847	0	0	0	1	0	1
Simplocaria semistriata (Fabricius, 1794)	8	8	39	14	1	70
Morychus aeneus (Fabricius, 1775)	1	0	0	0	2	3
Cytilus sericeus (Forster, 1771)	3	15	31	14	3	66
Byrrhus arietinus Steffahn, 1842	0	1	4	1	0	6
Byrrhus fasciatus (Forster, 1770)	2	7	15	0	3	27
Byrrhus pilula (Linnaeus, 1758)	6	15	45	16	2	84
Byrrhus pustulatus (Forster, 1770)	2	5	7	1	1	16
Porcinolus murinus (Fabricius, 1794)	1	1	1	0	0	3
Curimopsis nigrita (Palm, 1934)	0	0	11	0	0	11
Curimopsis setigera (Illiger, 1798)	3	0	0	0	0	3

Elmidae [12/8]

<i>Elmis aenea</i> (Müller, P.W.J., 1806)	4	15	17	18	19	73
<i>Esolus parallelepipedus</i> (Müller, P.W.J., 1806)	0	10	3	9	7	29
<i>Limnius volckmari</i> (Panzer, 1793)	0	12	6	18	14	50
<i>Normandia nitens</i> (Müller, P.W.J., 1817)	0	1	0	0	0	1
<i>Oulimnius troglodytes</i> (Gyllenhal, 1827)	0	0	0	5	0	5
<i>Oulimnius tuberculatus</i> (Müller, P.W.J., 1806)	1	11	6	16	5	39
<i>Riolus cupreus</i> (Müller, P.W.J., 1806)	0	3	1	1	1	6
<i>Riolus subviolaceus</i> (Müller, P.W.J., 1817)	0	5	0	2	1	8

Dryopidae [9/6]

<i>Pomatinus substriatus</i> (Müller, P.W.J., 1806)	0	2	1	0	0	3
<i>Dryops</i> (<i>Dryops</i>) <i>auriculatus</i> (Fourcroy, 1785)	3	1	2	1	2	9
<i>Dryops</i> (<i>Dryops</i>) <i>ernesti</i> des Gozis, 1886	3	12	17	17	6	55
<i>Dryops</i> (<i>Dryops</i>) <i>griseus</i> (Erichson, 1847)	0	0	0	1	0	1
<i>Dryops</i> (<i>Dryops</i>) <i>luridus</i> (Erichson, 1847)	5	1	8	3	1	18
<i>Dryops</i> (<i>Yrdops</i>) <i>nitidulus</i> (Heer, 1841)	0	0	1	2	3	6

Limnichidae [1/1]

<i>Limnichus pygmaeus</i> (Sturm, 1807)	0	1	6	0	0	7
---	---	---	---	---	---	---

Heteroceridae [9/5]

<i>Augyles maritimus</i> (Guérin-Méneville, 1844)	7	0	3	1	0	11
<i>Heterocerus fenestratus</i> (Thunberg, 1784)	18	2	5	5	3	33
<i>Heterocerus flexuosus</i> Stephens, 1828	3	1	4	1	0	9
<i>Heterocerus fuscus</i> Kiesenwetter, 1843	0	0	0	1	0	1
<i>Heterocerus marginatus</i> (Fabricius, 1787)	8	6	7	8	0	29

Psephenidae [1/1]

<i>Eubria palustris</i> Germar, 1818	0	7	1	4	0	12
--------------------------------------	---	---	---	---	---	----

Eucnemidae [6/4]

<i>Melasis buprestoides</i> (Linnaeus, 1761)	0	2	27	13	0	42
<i>Epiphanis cornutus</i> Eschscholtz, 1829	0	1	0	0	0	1
<i>Microrhagus pygmaeus</i> (Fabricius, 1792)	0	4	0	0	0	4
<i>Eucnemis capucina</i> Ahrens, 1812	0	0	1	0	0	1

Throscidae [5/3]

<i>Trixagus carinifrons</i> (de Bonvouloir, 1859)	1	1	15	1	0	18
<i>Trixagus dermestoides</i> (Linnaeus, 1767)	3	17	56	9	2	87
<i>Trixagus obtusus</i> (Curtis, 1827)	0	0	2	0	2	4

Elateridae [73/38]

<i>Agrypnus murinus</i> (Linnaeus, 1758)	20	3	4	1	0	28
<i>Hypnoidus riparius</i> (Fabricius, 1792)	11	16	79	23	24	153
<i>Actenicerus sjaelandicus</i> (Müller, O.F., 1764)	1	7	2	8	0	18
<i>Anostirus castaneus</i> (Linnaeus, 1758)	0	0	0	6	0	6
<i>Ctenicera cuprea</i> (Fabricius, 1775)	1	19	69	21	13	123
<i>Ctenicera pectinicornis</i> (Linnaeus, 1758)	3	8	13	6	3	33
<i>Aplotarsus incanus</i> (Gyllenhal, 1827)	25	47	87	21	14	194

Yorkshire Coleoptera checklist 2011

Paraphotistus impressus (Fabricius, 1792)	0	2	0	3	0	5
Paraphotistus nigricornis (Panzer, 1799)	24	2	2	1	0	29
Prosternon tessellatum (Linnaeus, 1758)	8	6	28	1	0	43
Selatosomus aeneus (Linnaeus, 1758)	4	2	12	7	1	26
Cidnopus aeruginosus (Olivier, 1790)	7	4	4	3	2	20
Kibunea minuta (Linnaeus, 1758)	24	15	93	4	2	138
Denticollis linearis (Linnaeus, 1758)	9	29	110	22	13	183
Athous (Athous) haemorrhoidalis (Fabricius, 1801)	58	72	547	60	19	756
Athous (Athous) vittatus (Fabricius, 1792)	4	13	23	7	3	50
Athous (Exanathrotus) subfuscus (Müller, O.F., 1764)	0	0	0	1	0	1
Athous (Orthathous) bicolor (Goeze, 1777)	12	3	32	3	0	50
Athous (Orthathous) campyloides Newman, 1833	0	0	1	2	0	3
Stenagostus rhombeus (Olivier, 1790)	0	0	0	1	0	1
Hemicrepidius hirtus (Herbst, 1784)	23	36	62	18	1	140
Adrastus pallens (Fabricius, 1792)	18	22	78	16	10	144
Agriotes acuminatus (Stephens, 1830)	24	43	155	27	7	256
Agriotes lineatus (Linnaeus, 1767)	21	12	7	3	0	43
Agriotes obscurus (Linnaeus, 1758)	44	36	158	39	13	290
Agriotes pallidulus (Illiger, 1807)	32	86	508	80	31	737
Agriotes sputator (Linnaeus, 1758)	21	14	23	6	3	67
Dalopius marginatus (Linnaeus, 1758)	14	82	232	56	12	396
Ampedus balteatus (Linnaeus, 1758)	9	18	41	3	2	73
Ampedus nigerrimus (Boisduval & Lacordaire, 1835)	0	0	1	0	0	1
Ampedus nigrinus (Herbst, 1784)	0	1	1	1	0	3
Ampedus pomorum (Herbst, 1784)	0	4	1	0	0	5
Ischnodes sanguinicollis (Panzer, 1793)	0	1	0	0	0	1
Sericus brunneus (Linnaeus, 1758)	2	4	12	5	1	24
Melanotus villosus (Geoffroy in Fourcroy, 1785)	12	22	59	19	4	116
Flautiauxellus maritimus (Curtis, 1840)	1	0	0	3	6	10
Oedostethus quadripustulatus (Fabricius, 1792)	12	4	10	1	1	28
Zorochochroa minimus (Boisduval & Lacordaire, 1835)	0	4	12	15	22	53

Lycidae [4/2]

Pyropterus nigroruber (De Geer, 1774)	2	5	25	2	0	34
Platycis minutus (Fabricius, 1787)	0	15	7	5	0	27

Lampyridae [3/1]

Lampyris noctiluca (Linnaeus, 1758)	6	11	94	16	6	133
-------------------------------------	---	----	----	----	---	-----

Cantharidae [41/38]

Podabrus alpinus (Paykull, 1798)	2	14	29	13	7	65
Ancistronycha abdominalis (Fabricius, 1798)	2	12	14	3	0	31
Cantharis cryptica Ashe, 1947	32	34	295	43	24	428
Cantharis decipiens Baudi, 1871	16	26	206	28	8	284
Cantharis figurata Mannerheim, 1843	0	0	7	3	1	11
Cantharis fusca Linnaeus, 1758	2	0	4	3	1	10
Cantharis lateralis Linnaeus, 1758	31	11	80	12	1	135
Cantharis livida Linnaeus, 1758	11	14	102	13	4	144
Cantharis nigra (De Geer, 1774)	45	30	187	31	2	295
Cantharis nigricans (Müller, O.F., 1776)	32	38	295	41	21	427

Yorkshire Coleoptera checklist 2011

<i>Cantharis pallida</i> Goeze, 1777	5	9	26	8	2	50
<i>Cantharis paludosa</i> Fallén, 1807	1	14	20	12	7	54
<i>Cantharis pellucida</i> Fabricius, 1792	9	32	139	17	12	209
<i>Cantharis rufa</i> Linnaeus, 1758	33	19	133	19	2	206
<i>Cantharis rustica</i> Fallén, 1807	10	13	59	10	6	98
<i>Cantharis thoracica</i> (Olivier, 1790)	18	6	25	8	2	59
<i>Rhagonycha elongata</i> (Fallén, 1807)	0	0	1	0	0	1
<i>Rhagonycha fulva</i> (Scopoli, 1763)	79	51	753	57	19	959
<i>Rhagonycha lignosa</i> (Müller, O.F., 1764)	8	29	154	15	6	212
<i>Rhagonycha limbata</i> Thomson, C.G., 1864	36	35	242	31	9	353
<i>Rhagonycha lutea</i> (Müller, O.F., 1764)	2	6	0	1	0	9
<i>Rhagonycha testacea</i> (Linnaeus, 1758)	5	23	64	26	12	130
<i>Rhagonycha translucida</i> (Krynicky, 1832)	0	8	5	1	0	14
<i>Crudosilis ruficollis</i> (Fabricius, 1775)	3	0	4	0	0	7
<i>Malthinus flaveolus</i> (Herbst, 1786)	10	21	97	11	2	141
<i>Malthinus frontalis</i> (Marsham, 1802)	2	6	12	8	1	29
<i>Malthinus seriepunctatus</i> Kiesenwetter, 1852	3	6	9	0	0	18
<i>Malthodes crassicornis</i> (Mäklin, 1846)	0	2	0	1	0	3
<i>Malthodes dispar</i> (Germar, 1824)	7	3	3	12	4	29
<i>Malthodes fibulatus</i> Kiesenwetter, 1852	0	4	0	3	0	7
<i>Malthodes flavoguttatus</i> Kiesenwetter, 1852	0	7	6	4	0	17
<i>Malthodes fuscus</i> (Waltl, 1838)	2	6	15	7	0	30
<i>Malthodes guttifer</i> Kiesenwetter, 1852	0	4	4	2	2	12
<i>Malthodes marginatus</i> (Latreille, 1806)	2	22	47	16	8	95
<i>Malthodes maurus</i> (Laporte, 1840)	0	4	0	1	0	5
<i>Malthodes minimus</i> (Linnaeus, 1758)	29	22	48	8	3	110
<i>Malthodes mysticus</i> Kiesenwetter, 1852	0	33	24	13	7	77
<i>Malthodes pumilus</i> (Brébisson, 1835)	6	9	24	7	1	47

Derodontidae [1/1]

<i>Laricobius erichsonii</i> Rosenhauer, 1846	0	2	0	0	0	2
---	---	---	---	---	---	---

Dermestidae [40/19]

<i>Dermestes</i> (<i>Dermestinus</i>) <i>frischii</i> Kugelann, 1792	0	2	1	0	0	3
<i>Dermestes</i> (<i>Dermestinus</i>) <i>maculatus</i> De Geer, 1774	3	3	6	3	0	15
<i>Dermestes</i> (<i>Dermestinus</i>) <i>murinus</i> Linnaeus, 1758	7	2	5	1	0	15
<i>Dermestes</i> (<i>Dermestinus</i>) <i>undulatus</i> Brahm, 1790	2	1	0	0	0	3
<i>Dermestes</i> (<i>Dermestes</i>) <i>ater</i> De Geer, 1774	0	0	1	0	0	1
<i>Dermestes</i> (<i>Dermestes</i>) <i>haemorrhoidalis</i> Küster, 1852	0	0	1	0	0	1
<i>Dermestes</i> (<i>Dermestes</i>) <i>lardarius</i> Linnaeus, 1758	4	5	15	1	0	25
<i>Dermestes</i> (<i>Dermestes</i>) <i>peruvianus</i> Laporte de Castelnau, 1840	0	0	7	0	0	7
<i>Attagenus pelli</i> (Linnaeus, 1758)	9	4	30	3	2	48
<i>Attagenus unicolor</i> (Brahm, 1791)	0	0	0	1	0	1
<i>Trogoderma granarium</i> Everts, 1898	0	1	0	0	0	1
<i>Trogoderma inclusum</i> LeConte, 1854	0	0	1	0	0	1
<i>Reesa vespulae</i> (Milliron, 1939)	2	0	0	0	0	2
<i>Megatoma undata</i> (Linnaeus, 1758)	5	2	3	4	0	14
<i>Ctesias serra</i> (Fabricius, 1792)	0	9	3	4	0	16
<i>Anthrenus</i> (<i>Florilinus</i>) <i>museorum</i> (Linnaeus, 1761)	1	3	6	0	0	10
<i>Anthrenus</i> (<i>Helocerus</i>) <i>fuscus</i> Olivier, 1789	6	9	36	5	0	56

Yorkshire Coleoptera checklist 2011

<i>Anthrenus</i> (<i>Nathrenus</i>) <i>verbasci</i> (Linnaeus, 1767)	6	3	22	5	0	36
<i>Trinodes</i> <i>hirtus</i> (Fabricius, 1781)	0	0	1	0	0	1
<u>Bostrichidae [5/5]</u>						
<i>Bostrichus</i> <i>capucinus</i> (Linnaeus, 1758)	1	0	2	1	0	4
<i>Rhyzopertha</i> <i>dominica</i> (Fabricius, 1792)	2	1	2	0	0	5
<i>Lyctus</i> <i>brunneus</i> (Stephens, 1830)	0	3	6	0	0	9
<i>Lyctus</i> <i>linearis</i> (Goeze, 1777)	0	3	1	0	0	4
<i>Lyctus</i> <i>planicollis</i> LeConte, 1858	0	0	1	0	0	1
<u>Anobiidae [53/38]</u>						
<i>Hedobia</i> (<i>Ptinomorphus</i>) <i>imperialis</i> (Linnaeus, 1767)	8	6	12	6	0	32
<i>Gibbium</i> <i>aequinoctiale</i> Boieldieu, 1854	0	0	6	0	0	6
<i>Gibbium</i> <i>psylloides</i> (de Czenpinski, 1778)	1	1	7	0	0	9
<i>Stethomezium</i> <i>squamosum</i> Hinton, 1943	0	0	1	0	0	1
<i>Mezium</i> <i>affine</i> Boieldieu, 1856	0	1	1	1	0	3
<i>Trigonogenius</i> <i>globulus</i> Solier, 1849	1	2	1	1	0	5
<i>Niptus</i> <i>hololeucus</i> (Faldermann, 1835)	4	5	14	5	0	28
<i>Tipnus</i> <i>unicolor</i> (Piller & Mitterpacher, 1783)	4	3	6	2	0	15
<i>Pseudeurostus</i> <i>hilleri</i> (Reitter, 1877)	0	0	1	0	0	1
<i>Ptinus</i> <i>clavipes</i> Panzer, 1792	0	0	1	0	0	1
<i>Ptinus</i> <i>fur</i> (Linnaeus, 1758)	12	5	11	1	0	29
<i>Ptinus</i> <i>lichenum</i> Marsham, 1802	0	0	2	0	0	2
<i>Ptinus</i> <i>palliatu</i> s Perris, 1847	0	0	1	0	0	1
<i>Ptinus</i> <i>pusillus</i> Sturm, 1837	0	0	1	0	0	1
<i>Ptinus</i> <i>sexpunctatus</i> Panzer, 1792	0	0	3	0	0	3
<i>Ptinus</i> <i>subpilosus</i> Sturm, 1837	0	0	2	0	0	2
<i>Ptinus</i> <i>tectus</i> Boieldieu, 1856	5	5	20	3	0	33
<i>Grynobius</i> <i>planus</i> (Fabricius, 1787)	4	17	9	6	1	37
<i>Dryophilus</i> <i>pusillus</i> (Gyllenhal, 1808)	5	22	14	3	5	49
<i>Ochina</i> <i>ptinoides</i> (Marsham, 1802)	0	2	2	3	0	7
<i>Xestobium</i> <i>rufovillosum</i> (De Geer, 1774)	5	8	6	6	1	26
<i>Ernobius</i> <i>mollis</i> (Linnaeus, 1758)	5	8	13	3	2	31
<i>Ernobius</i> <i>nigrinus</i> (Sturm, 1837)	1	2	0	0	0	3
<i>Stegobium</i> <i>paniceum</i> (Linnaeus, 1758)	2	4	28	3	0	37
<i>Gastrallus</i> <i>immarginatus</i> (Müller, P.W.J., 1821)	0	0	1	0	0	1
<i>Anobium</i> Fabricius, 1775	0	0	1	0	0	1
<i>Anobium</i> <i>fulvicorne</i> (Sturm, 1837)	6	5	18	8	0	37
<i>Anobium</i> <i>inexpectatum</i> Lohse, 1954	1	0	4	1	1	7
<i>Anobium</i> <i>punctatum</i> (De Geer, 1774)	16	12	30	13	1	72
<i>Hadrobregmus</i> <i>denticollis</i> (Creutzer in Panzer, 1796)	0	0	2	0	0	2
<i>Ptilinus</i> <i>pectinicornis</i> (Linnaeus, 1758)	17	32	72	20	3	144
<i>Lasioderma</i> <i>serricorne</i> (Fabricius, 1792)	0	0	2	0	0	2
<i>Dorcatoma</i> <i>chrysomelina</i> (Sturm, 1837)	1	3	4	9	0	17
<i>Dorcatoma</i> <i>dresdensis</i> Herbst, 1792	4	3	2	0	0	9
<i>Dorcatoma</i> <i>flavicornis</i> (Fabricius, 1792)	1	4	4	5	0	14
<i>Dorcatoma</i> <i>substriata</i> Hummel, 1829	0	4	1	0	0	5
<i>Caenocara</i> <i>bovistae</i> (Hoffmann, J.J., 1803)	0	0	1	0	0	1
<i>Anitys</i> <i>rubens</i> (Hoffmann, J.J., 1803)	0	1	1	3	0	5

Lymexilidae [2/1]

Hylecoetus dermestoides (Linnaeus, 1761)	5	17	31	7	0	60
--	---	----	----	---	---	----

Phloiophilidae [1/1]

Phloiophilus edwardsii Stephens, 1830	1	5	0	1	0	7
---------------------------------------	---	---	---	---	---	---

Trogossitidae [5/4]

Thymalus limbatus (Fabricius, 1787)	0	5	2	6	0	13
Lophocateres pusillus (Klug, 1832)	0	0	1	0	0	1
Tenebroides mauritanicus (Linnaeus, 1758)	1	2	4	0	0	7
Nemozoma elongatum (Linnaeus, 1761)	1	0	1	0	0	2

Cleridae [14/8]

Tillus elongatus (Linnaeus, 1758)	0	3	0	0	0	3
Opilo mollis (Linnaeus, 1758)	0	1	0	0	0	1
Thanasimus formicarius (Linnaeus, 1758)	9	7	8	2	0	26
Tarsostenus univittatus (Rossi, 1792)	0	0	1	0	0	1
Korynetes caeruleus (De Geer, 1775)	2	1	2	1	0	6
Necrobia ruficollis (Fabricius, 1775)	3	3	3	2	0	11
Necrobia rufipes (De Geer, 1775)	8	5	9	1	0	23
Necrobia violacea (Linnaeus, 1758)	12	4	17	5	2	40

Dasytidae [9/8]

Aplocnemus impressus (Marsham, 1802)	0	1	2	0	0	3
Aplocnemus nigricornis (Fabricius, 1792)	0	6	0	1	0	7
Dasytes aeratus Stephens, 1830	4	15	7	9	3	38
Dasytes niger (Linnaeus, 1761)	0	0	1	0	0	1
Dasytes plumbeus (Müller, O.F., 1776)	0	2	3	0	0	5
Dasytes virens (Marsham, 1802)	0	0	2	0	0	2
Psilothrix viridicoeruleus (Geoffroy in Fourcroy, 1785)	0	0	1	0	0	1
Dolichosoma lineare (Rossi, 1794)	4	0	0	0	0	4

Malachiidae [16/4]

Ebaeus pedicularius (Linnaeus, 1758)	0	0	1	0	0	1
Cordylepherus viridis (Fabricius, 1787)	1	0	1	2	0	4
Malachius bipustulatus (Linnaeus, 1758)	27	31	210	32	8	308
Anthocomus rufus (Herbst, 1784)	5	0	42	0	0	47

Sphindidae [2/2]

Sphindus dubius (Gyllenhal, 1808)	0	7	1	0	0	8
Aspidiphorus orbiculatus (Gyllenhal, 1808)	2	9	18	2	0	31

Kateretidae [9/8]

Kateretes pedicularius (Linnaeus, 1758)	1	3	4	7	0	15
Kateretes pusillus (Thunberg, 1794)	13	9	1	6	2	31
Kateretes rufilabris (Latreille, 1807)	26	10	31	15	6	88
Brachypterus glaber (Stephens, 1832)	57	31	230	34	12	364
Brachypterus urticae (Fabricius, 1792)	68	45	298	42	20	473
Brachypterolus linariae (Stephens, 1830)	0	0	4	0	0	4
Brachypterolus pulicarius (Linnaeus, 1758)	4	0	28	2	1	35

Yorkshire Coleoptera checklist 2011

<i>Brachypterolus vestitus</i> (Kiesenwetter, 1850)	0	0	2	0	0	2
<u>Nitidulidae [88/68]</u>						
<i>Carpophilus dimidiatus</i> (Fabricius, 1792)	1	0	3	0	0	4
<i>Carpophilus hemipterus</i> (Linnaeus, 1758)	1	1	3	2	0	7
<i>Carpophilus ligneus</i> Murray, 1864	0	1	4	1	0	6
<i>Carpophilus marginellus</i> Motschulsky, 1858	6	0	28	1	1	36
<i>Carpophilus mutilatus</i> Erichson, 1843	0	0	3	0	0	3
<i>Carpophilus obsoletus</i> Erichson, 1843	0	0	2	0	0	2
<i>Carpophilus sexpustulatus</i> (Fabricius, 1792)	1	1	15	4	0	21
<i>Epuraea</i> (<i>Epuraea</i>) <i>aestiva</i> (Linnaeus, 1758)	30	42	153	26	13	264
<i>Epuraea</i> (<i>Epuraea</i>) <i>angustula</i> Sturm, 1844	0	2	3	1	0	6
<i>Epuraea</i> (<i>Epuraea</i>) <i>biguttata</i> (Thunberg, 1784)	13	22	85	17	5	142
<i>Epuraea</i> (<i>Epuraea</i>) <i>fuscicollis</i> (Stephens, 1832)	0	1	0	2	0	3
<i>Epuraea</i> (<i>Epuraea</i>) <i>guttata</i> (Olivier, 1811)	0	0	2	0	0	2
<i>Epuraea</i> (<i>Epuraea</i>) <i>longula</i> Erichson, 1845	0	1	0	1	0	2
<i>Epuraea</i> (<i>Epuraea</i>) <i>marseuli</i> Reitter, 1872	3	4	9	3	0	19
<i>Epuraea</i> (<i>Epuraea</i>) <i>melina</i> Erichson, 1843	4	7	11	1	0	23
<i>Epuraea</i> (<i>Epuraea</i>) <i>pallescens</i> (Stephens, 1832)	5	5	12	2	0	24
<i>Epuraea</i> (<i>Epuraea</i>) <i>rufomarginata</i> (Stephens, 1830)	0	3	8	0	0	11
<i>Epuraea</i> (<i>Epuraea</i>) <i>silacea</i> (Herbst, 1783)	3	13	13	8	1	38
<i>Epuraea</i> (<i>Epuraea</i>) <i>terminalis</i> Mannerheim, 1843	0	3	1	0	0	4
<i>Epuraea</i> (<i>Epuraea</i>) <i>thoracica</i> Tournier, 1872	0	3	0	1	0	4
<i>Epuraea</i> (<i>Epuraea</i>) <i>variegata</i> (Herbst, 1793)	0	1	0	0	0	1
<i>Epuraea</i> (<i>Epuraeanella</i>) <i>limbata</i> (Fabricius, 1787)	2	1	1	0	0	4
<i>Epuraea</i> (<i>Micruria</i>) <i>melanocephala</i> (Marsham, 1802)	17	22	140	15	5	199
<i>Pria dulcamarae</i> (Scopoli, 1763)	7	1	18	6	0	32
<i>Meligethes aeneus</i> (Fabricius, 1775)	169	129	992	112	56	1458
<i>Meligethes atramentarius</i> Förster, 1849	0	1	0	1	0	2
<i>Meligethes atratus</i> (Olivier, 1790)	7	12	60	10	0	89
<i>Meligethes bidens</i> Brisout de Barneville, 1863	1	0	0	0	0	1
<i>Meligethes brevis</i> Sturm, 1845	4	5	0	1	0	10
<i>Meligethes brunnicornis</i> Sturm, 1845	3	6	10	1	0	20
<i>Meligethes carinulatus</i> Förster, 1849	19	9	28	1	2	59
<i>Meligethes difficilis</i> (Heer, 1841)	12	4	25	0	0	41
<i>Meligethes flavimanus</i> Stephens, 1830	4	10	3	3	2	22
<i>Meligethes fulvipes</i> Brisout de Barneville, 1863	5	3	1	1	0	10
<i>Meligethes gagathinus</i> Erichson, 1845	0	1	0	0	0	1
<i>Meligethes incanus</i> Sturm, 1845	0	0	1	0	0	1
<i>Meligethes kunzei</i> Erichson, 1845	0	0	0	1	0	1
<i>Meligethes lugubris</i> Sturm, 1845	2	0	1	0	0	3
<i>Meligethes morosus</i> Erichson, 1845	11	4	11	0	1	27
<i>Meligethes nigrescens</i> Stephens, 1830	35	14	67	9	1	126
<i>Meligethes obscurus</i> Erichson, 1845	2	7	8	4	0	21
<i>Meligethes ochropus</i> Sturm, 1845	0	0	2	0	0	2
<i>Meligethes ovatus</i> Sturm, 1845	4	2	5	0	0	11
<i>Meligethes pedicularius</i> (Gyllenhal, 1808)	7	2	14	1	2	26
<i>Meligethes planiusculus</i> (Heer, 1841)	0	1	8	0	0	9
<i>Meligethes rotundicollis</i> Brisout de Barneville, 1863	0	0	1	0	0	1
<i>Meligethes ruficornis</i> (Marsham, 1802)	1	0	0	0	0	1

Yorkshire Coleoptera checklist 2011

Meligethes solidus (Kugelann, 1794)	0	0	2	3	0	5
Meligethes subrugosus (Gyllenhal, 1808)	7	0	0	0	0	7
Meligethes viridescens (Fabricius, 1787)	4	14	45	6	2	71
Nitidula bipunctata (Linnaeus, 1758)	4	5	9	5	2	25
Nitidula flavomaculata Rossi, 1790	0	1	0	0	0	1
Nitidula rufipes (Linnaeus, 1767)	1	0	8	3	0	12
Omosita colon (Linnaeus, 1758)	12	1	15	6	1	35
Omosita depressa (Linnaeus, 1758)	0	3	12	3	0	18
Omosita discoidea (Fabricius, 1775)	11	7	35	13	1	67
Soronia grisea (Linnaeus, 1758)	7	8	24	9	1	49
Soronia punctatissima (Illiger, 1794)	0	4	5	6	0	15
Cychramus luteus (Fabricius, 1787)	1	11	13	2	0	27
Pocadius adustus Reitter, 1888	0	4	8	0	0	12
Pocadius ferrugineus (Fabricius, 1775)	3	3	13	2	1	22
Thalycra fervida (Olivier, 1790)	0	0	1	2	0	3
Cryptarcha strigata (Fabricius, 1787)	0	0	2	2	0	4
Cryptarcha undata (Olivier, 1790)	0	0	0	3	0	3
Glischrochilus (Glischrochilus) quadripunctatus (Linnaeus, 1758)	4	8	7	2	1	22
Glischrochilus (Librodor) hortensis (Geoffroy in Fourcroy, 1785)	28	28	81	34	3	174
Glischrochilus (Librodor) quadriguttatus (Fabricius, 1777)	1	3	5	3	0	12
Pityophagus ferrugineus (Linnaeus, 1761)	0	2	1	0	0	3

Monotomidae [23/20]

Rhizophagus (Eurhizophagus) depressus (Fabricius, 1792)	1	3	11	3	0	18
Rhizophagus (Rhizophagus) bipustulatus (Fabricius, 1792)	10	16	74	23	2	125
Rhizophagus (Rhizophagus) dispar (Paykull, 1800)	12	24	183	34	2	255
Rhizophagus (Rhizophagus) ferrugineus (Paykull, 1800)	2	6	5	2	0	15
Rhizophagus (Rhizophagus) nitidulus (Fabricius, 1798)	0	3	24	6	0	33
Rhizophagus (Rhizophagus) oblongicollis Blatch & Horner, 1892	0	1	0	0	0	1
Rhizophagus (Rhizophagus) parallelocolis Gyllenhal, 1827	1	4	6	0	0	11
Rhizophagus (Rhizophagus) perforatus Erichson, 1845	3	4	28	7	0	42
Rhizophagus (Rhizophagus) picipes (Olivier, 1790)	1	0	7	3	0	11
Rhizophagus (Anomophagus) cribratus Gyllenhal, 1827	0	0	4	4	0	8
Cyanostolus aeneus (Richter, 1820)	0	1	0	0	0	1
Monotoma (Gyrocecis) angusticollis (Gyllenhal, 1827)	0	3	0	0	0	3
Monotoma (Gyrocecis) conicicollis Guérin-Ménéville, 1837	0	4	6	0	0	10
Monotoma (Gyrocecis) quadrifoveolata Aubé, 1838	0	1	0	0	0	1
Monotoma (Monotoma) bicolor Villa & Villa, 1835	5	2	27	5	1	40
Monotoma (Monotoma) brevicollis Aubé, 1838	0	0	1	0	0	1
Monotoma (Monotoma) longicollis (Gyllenhal, 1827)	8	7	14	3	0	32
Monotoma (Monotoma) picipes Herbst, 1793	12	8	33	9	2	64
Monotoma (Monotoma) spinicollis Aubé, 1838	4	0	6	3	0	13
Monotoma (Monotoma) testacea Motschulsky, 1845	0	0	6	2	0	8

Silvanidae [12/9]

Uleiota planata (Linnaeus, 1761)	6	0	0	0	0	6
Cryptomorpha desjardinsii (Guérin-Ménéville, 1844)	0	0	1	0	0	1
Psammoecus bipunctatus (Fabricius, 1792)	3	0	5	0	0	8
Nausibius clavicornis (Kugelann, 1794)	0	0	1	0	0	1
Ahasverus advena (Waltl, 1834)	8	2	18	3	0	31

Yorkshire Coleoptera checklist 2011

<i>Oryzaephilus mercator</i> (Fauvel, 1889)	1	0	1	0	0	2
<i>Oryzaephilus surinamensis</i> (Linnaeus, 1758)	2	0	3	0	0	5
<i>Silvanus bidentatus</i> (Fabricius, 1792)	0	0	1	0	0	1
<i>Silvanus unidentatus</i> (Olivier, 1790)	0	2	2	0	0	4

Cucujidae [2/2]

<i>Pediacus depressus</i> (Herbst, 1797)	0	0	1	0	0	1
<i>Pediacus dermestoides</i> (Fabricius, 1792)	0	9	4	2	1	16

Laemophloeidae [11/2]

<i>Cryptolestes ferrugineus</i> (Stephens, 1831)	3	5	16	3	0	27
<i>Cryptolestes pusillus</i> (Schönherr, 1817)	0	0	3	0	0	3

Phalacridae [15/11]

<i>Phalacrus caricis</i> Sturm, 1807	5	4	8	1	0	18
<i>Phalacrus corruscus</i> (Panzer, 1796)	1	0	4	0	0	5
<i>Phalacrus fimetarius</i> (Fabricius, 1775)	21	2	18	3	0	44
<i>Phalacrus substriatus</i> Gyllenhal, 1813	0	1	2	0	0	3
<i>Olibrus aeneus</i> (Fabricius, 1792)	33	5	63	10	1	112
<i>Olibrus affinis</i> (Sturm, 1807)	1	0	1	0	0	2
<i>Olibrus corticalis</i> (Panzer, 1796)	6	0	2	0	0	8
<i>Olibrus liquidus</i> Erichson, 1845	5	0	6	1	0	12
<i>Olibrus pygmaeus</i> (Sturm, 1807)	1	0	0	0	0	1
<i>Stilbus oblongus</i> (Erichson, 1845)	4	0	9	0	0	13
<i>Stilbus testaceus</i> (Panzer, 1796)	111	40	158	26	8	343

Cryptophagidae [103/73]

<i>Telmatophilus caricis</i> (Olivier, 1790)	6	7	40	7	0	60
<i>Telmatophilus schoenherrii</i> (Gyllenhal, 1808)	0	0	3	0	0	3
<i>Telmatophilus typhae</i> (Fallén, 1802)	22	5	77	10	3	117
<i>Paramecosoma melanocephalum</i> (Herbst, 1793)	0	6	5	8	2	21
<i>Henoticus serratus</i> (Gyllenhal, 1808)	2	7	11	3	0	23
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>acutangulus</i> Gyllenhal, 1827	5	5	19	3	0	32
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>angustus</i> Ganglbauer, 1899	1	0	0	0	0	1
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>badius</i> Sturm, 1845	1	1	0	4	0	6
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>cellaris</i> (Scopoli, 1763)	1	4	3	1	0	9
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>corticinus</i> Thomson, C.G., 1863	0	0	1	0	0	1
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>dentatus</i> (Herbst, 1793)	8	9	58	12	4	91
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>distinguendus</i> Sturm, 1845	9	1	18	7	1	36
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>immixtus</i> Rey, 1889	1	3	2	2	0	8
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>labilis</i> Erichson, 1846	0	0	1	0	0	1
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>lapponicus</i> Gyllenhal, 1827	0	1	0	0	0	1
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>laticollis</i> Lucas, 1846	7	6	19	2	4	38
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>lycoperdi</i> (Scopoli, 1763)	5	1	31	5	0	42
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>obsoletus</i> Reitter, 1879	0	0	0	1	0	1
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>pallidus</i> Sturm, 1845	4	4	4	1	0	13
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>pilosus</i> Gyllenhal, 1827	9	4	43	7	0	63
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>populi</i> Paykull, 1800	0	1	5	1	0	7
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>pseudodentatus</i> Bruce, 1934	3	3	20	9	3	38
<i>Cryptophagus</i> (<i>Cryptophagus</i>) <i>pubescens</i> Sturm, 1845	4	5	12	3	2	26

Yorkshire Coleoptera checklist 2011

Cryptophagus (Cryptophagus) ruficornis Stephens, 1830	5	0	13	1	0	19
Cryptophagus (Cryptophagus) saginatus Sturm, 1845	2	1	3	0	0	6
Cryptophagus (Cryptophagus) scanicus (Linnaeus, 1758)	3	12	23	8	2	48
Cryptophagus (Cryptophagus) schmidtii Leunis in Sturm, 1845	0	1	3	0	0	4
Cryptophagus (Cryptophagus) scutellatus Newman, 1834	5	3	23	3	0	34
Cryptophagus (Cryptophagus) setulosus Sturm, 1845	8	13	15	4	4	44
Cryptophagus (Cryptophagus) subdepressus Gyllenhal, 1827	0	1	0	0	0	1
Micrambe abietis (Paykull, 1798)	1	0	0	0	0	1
Micrambe lindbergorum (Bruce, 1934)	6	0	0	0	1	7
Micrambe villosa (Heer, 1841)	13	4	10	0	0	27
Micrambe vini (Panzer, 1797)	48	18	84	5	15	170
Antherophagus nigricornis (Fabricius, 1787)	4	7	33	4	1	49
Antherophagus pallens (Linnaeus, 1758)	2	5	23	8	1	39
Caenoscelis ferruginea (Sahlberg, C.R., 1820)	0	3	4	0	0	7
Caenoscelis sibirica Reitter, 1889	0	0	1	0	0	1
Caenoscelis subdeplanata Brisout de Barneville, 1882	2	0	19	1	0	22
Atomaria (Anchicera) apicalis Erichson, 1846	9	6	30	3	0	48
Atomaria (Anchicera) atra (Herbst, 1793)	1	1	0	0	0	2
Atomaria (Anchicera) atricapilla Stephens, 1830	28	21	128	16	6	199
Atomaria (Anchicera) basalis Erichson, 1846	8	1	2	1	0	12
Atomaria (Anchicera) fuscata (Schönherr, 1808)	56	29	143	34	4	266
Atomaria (Anchicera) fuscipes (Gyllenhal, 1808)	2	5	2	2	1	12
Atomaria (Anchicera) hislopi Wollaston, 1857	0	1	0	0	0	1
Atomaria (Anchicera) lewisi Reitter, 1877	67	31	251	41	10	400
Atomaria (Anchicera) mesomela (Herbst, 1792)	11	1	27	11	3	53
Atomaria (Anchicera) morio Kolenati, 1846	2	0	0	0	0	2
Atomaria (Anchicera) munda Erichson, 1846	2	1	7	3	0	13
Atomaria (Anchicera) nigripennis (Kugelann, 1794)	0	0	1	0	0	1
Atomaria (Anchicera) nitidula (Marsham, 1802)	64	33	267	39	12	415
Atomaria (Anchicera) peltata Kraatz, 1853	0	1	0	0	0	1
Atomaria (Anchicera) pusilla (Paykull, 1798)	3	7	7	4	2	23
Atomaria (Anchicera) rhenana Kraatz, 1853	11	0	0	0	0	11
Atomaria (Anchicera) rubella Heer, 1841	17	5	44	10	1	77
Atomaria (Anchicera) scutellaris Motschulsky, 1849	0	0	3	0	0	3
Atomaria (Anchicera) testacea Stephens, 1830	74	16	165	34	17	306
Atomaria (Atomaria) barani Brisout de Barneville, 1863	1	0	0	2	0	3
Atomaria (Atomaria) diluta Erichson, 1846	0	0	2	5	0	7
Atomaria (Atomaria) fimetarii (Fabricius, 1792)	3	1	5	2	0	11
Atomaria (Atomaria) impressa Erichson, 1846	0	1	0	1	1	3
Atomaria (Atomaria) linearis Stephens, 1830	20	16	50	11	7	104
Atomaria (Atomaria) lohsei Johnson & Strand, A., 1968	0	1	3	0	0	4
Atomaria (Atomaria) nigrirostris Stephens 1830	8	7	57	10	2	84
Atomaria (Atomaria) nigriventris Stephens, 1830	0	2	3	1	0	6
Atomaria (Atomaria) pulchra Erichson, 1846	0	5	4	1	0	10
Atomaria (Atomaria) puncticollis Thomson, C.G., 1868	1	0	0	0	0	1
Atomaria (Atomaria) punctithorax Reitter, 1887	1	0	0	0	0	1
Atomaria (Atomaria) strandi Johnson, 1967	0	1	3	0	0	4
Atomaria (Atomaria) umbrina (Gyllenhal, 1827)	0	1	1	0	0	2
Ootypus globosus (Waltl, 1838)	5	5	9	3	0	22
Ephistemus globulus (Paykull, 1798)	37	12	125	18	7	199

Erotylidae [7/6]

<i>Dacne bipustulata</i> (Thunberg, 1781)	8	14	40	22	2	86
<i>Dacne rufifrons</i> (Fabricius, 1775)	3	9	7	8	1	28
<i>Triplax aenea</i> (Schaller, 1783)	2	13	70	15	7	107
<i>Triplax russica</i> (Linnaeus, 1758)	2	0	15	0	0	17
<i>Triplax scutellaris</i> Charpentier, 1825	0	2	0	0	0	2
<i>Tritoma bipustulata</i> Fabricius, 1775	0	2	4	0	0	6

Byturidae [2/2]

<i>Byturus ochraceus</i> (Scriba, 1790)	10	9	52	9	1	81
<i>Byturus tomentosus</i> (De Geer, 1774)	29	54	280	27	10	400

Biphyllidae [2/1]

<i>Biphyllus lunatus</i> (Fabricius, 1787)	13	8	20	8	1	50
--	----	---	----	---	---	----

Bothrideridae [5/2]

<i>Teredus cylindricus</i> (Olivier, 1790)	0	0	2	0	0	2
<i>Anommatus duodecimstriatus</i> (Müller, P.W.J., 1821)	4	1	4	0	0	9

Cerylonidae [5/3]

<i>Cerylon fagi</i> Brisout de Barneville, 1867	1	0	0	1	0	2
<i>Cerylon ferrugineum</i> Stephens, 1830	1	17	74	19	4	115
<i>Cerylon histeroides</i> (Fabricius, 1792)	4	3	26	14	4	51

Alexiidae [1/1]

<i>Sphaerosoma pilosum</i> (Panzer, 1793)	0	0	1	0	0	1
---	---	---	---	---	---	---

Endomychidae [8/3]

<i>Endomychus coccineus</i> (Linnaeus, 1758)	4	12	20	11	0	47
<i>Symbiotes latus</i> Redtenbacher, 1849	0	0	1	0	0	1
<i>Mycetaea subterranea</i> (Fabricius, 1801)	27	5	33	3	1	69

Coccinellidae [53/41]

<i>Coccidula rufa</i> (Herbst, 1783)	96	46	325	46	11	524
<i>Coccidula scutellata</i> (Herbst, 1783)	10	2	33	2	0	47
<i>Rhyzobius litura</i> (Fabricius, 1787)	72	43	223	26	11	375
<i>Clitostethus arcuatus</i> (Rossi, 1794)	0	0	4	0	0	4
<i>Stethorus punctillum</i> (Weise, 1891)	1	1	4	0	0	6
<i>Scymnus</i> (<i>Scymnus</i>) <i>femoralis</i> (Gyllenhal, 1827)	3	2	3	0	0	8
<i>Scymnus</i> (<i>Scymnus</i>) <i>frontalis</i> (Fabricius, 1787)	2	1	0	0	0	3
<i>Scymnus</i> (<i>Scymnus</i>) <i>nigrinus</i> Kugelann, 1794	1	2	2	2	0	7
<i>Scymnus</i> (<i>Scymnus</i>) <i>schmidti</i> Fürsch, 1958	3	0	0	0	0	3
<i>Scymnus</i> (<i>Pullus</i>) <i>auritus</i> Thunberg, 1795	8	4	18	4	0	34
<i>Scymnus</i> (<i>Pullus</i>) <i>suturalis</i> Thunberg, 1795	12	19	19	3	2	55
<i>Scymnus</i> (<i>Neopullus</i>) <i>haemorrhoidalis</i> Herbst, 1797	3	3	12	1	0	19
<i>Scymnus</i> (<i>Neopullus</i>) <i>limbatus</i> Stephens, 1832	2	0	3	1	0	6
<i>Nephus redtenbacheri</i> (Mulsant, 1846)	12	7	16	5	0	40
<i>Hyperaspis pseudopustulata</i> Mulsant, 1853	2	4	5	0	0	11
<i>Chilocorus bipustulatus</i> (Linnaeus, 1758)	4	21	31	1	0	57
<i>Chilocorus renipustulatus</i> (Scriba, 1791)	21	11	85	18	2	137

Yorkshire Coleoptera checklist 2011

<i>Exochomus nigromaculatus</i> (Goeze, 1777)	0	0	2	0	0	2
<i>Exochomus quadripustulatus</i> (Linnaeus, 1758)	16	12	143	12	2	185
<i>Anisosticta novemdecimpunctata</i> (Linnaeus, 1758)	34	9	192	34	7	276
<i>Tytthaspis sedecimpunctata</i> (Linnaeus, 1758)	0	1	5	2	0	8
<i>Myzia oblongoguttata</i> (Linnaeus, 1758)	4	14	11	7	2	38
<i>Myrrha octodecimguttata</i> (Linnaeus, 1758)	12	13	20	10	0	55
<i>Propylea quattuordecimpunctata</i> (Linnaeus, 1758)	110	87	1086	78	31	1392
<i>Calvia quattuordecimguttata</i> (Linnaeus, 1758)	31	44	313	50	25	463
<i>Halyzia sedecimguttata</i> (Linnaeus, 1758)	6	7	63	9	3	88
<i>Psyllobora vigintiduopunctata</i> (Linnaeus, 1758)	63	15	398	30	5	511
<i>Anatis ocellata</i> (Linnaeus, 1758)	6	22	69	14	5	116
<i>Aphidecta oblitterata</i> (Linnaeus, 1758)	10	37	46	13	17	123
<i>Hippodamia</i> (<i>Hippodamia</i>) <i>tredecimpunctata</i> (Linnaeus, 1758)	1	2	3	0	0	6
<i>Hippodamia</i> (<i>Adonia</i>) <i>variegata</i> (Goeze, 1777)	22	5	95	13	1	136
<i>Coccinella hieroglyphica</i> Linnaeus, 1758	6	18	46	4	1	75
<i>Coccinella magnifica</i> Redtenbacher, 1843	0	0	5	0	0	5
<i>Coccinella quinquepunctata</i> Linnaeus, 1758	0	2	0	0	0	2
<i>Coccinella septempunctata</i> Linnaeus, 1758	124	85	1442	90	48	1789
<i>Coccinella undecimpunctata</i> Linnaeus, 1758	40	8	156	7	3	214
<i>Adalia bipunctata</i> (Linnaeus, 1758)	67	45	751	50	24	937
<i>Adalia decempunctata</i> (Linnaeus, 1758)	47	44	350	43	14	498
<i>Harmonia axyridis</i> (Pallas, 1773)	12	0	43	1	3	59
<i>Harmonia quadripunctata</i> (Pontoppidan, 1763)	2	1	33	2	0	38
<i>Subcoccinella vigintiquatuorpunctata</i> (Linnaeus, 1758)	48	10	28	1	0	87

Corylophidae [11/8]

<i>Sericoderus lateralis</i> (Gyllenhal, 1827)	10	1	24	2	2	39
<i>Corylophus cassidoides</i> (Marsham, 1802)	2	0	4	0	0	6
<i>Corylophus sublaevipennis</i> Jacquelin du Val, 1859	3	0	0	0	0	3
<i>Orthoperus aequalis</i> Sharp, 1885	0	0	1	0	0	1
<i>Orthoperus atomus</i> (Gyllenhal, 1808)	0	0	2	1	0	3
<i>Orthoperus brunnipes</i> (Gyllenhal, 1808)	9	2	0	0	0	11
<i>Orthoperus corticalis</i> (Redtenbacher, 1849)	1	1	4	0	0	6
<i>Orthoperus nigrescens</i> Stephens, 1829	1	0	2	0	0	3

Latridiidae [56/38]

<i>Lithostygnus serripennis</i> Broun, 1914	14	0	11	1	0	26
<i>Stephostethus angusticollis</i> (Gyllenhal, 1827)	1	0	3	0	0	4
<i>Stephostethus lardarius</i> (De Geer, 1775)	70	39	158	39	5	311
<i>Cartodere</i> (<i>Cartodere</i>) <i>constricta</i> (Gyllenhal 1827)	0	0	2	0	0	2
<i>Cartodere</i> (<i>Aridius</i>) <i>bifasciata</i> (Reitter, 1877)	111	42	350	47	9	559
<i>Cartodere</i> (<i>Aridius</i>) <i>nodifer</i> (Westwood, 1839)	76	38	272	44	9	439
<i>Cartodere</i> (<i>Aridius</i>) <i>norvegica</i> (Strand, 1940)	0	0	1	0	0	1
<i>Latridius anthracinus</i> Mannerheim, 1844	17	1	6	2	1	27
<i>Latridius assimilis</i> (Mannerheim, 1844)	2	2	16	3	1	24
<i>Latridius consimilis</i> Mannerheim, 1844	1	0	1	3	0	5
<i>Latridius minutus</i> (Linnaeus, 1767)	1	4	17	1	1	24
<i>Thes bergrothi</i> (Reitter, 1880)	0	2	2	2	0	6
<i>Enicmus brevicornis</i> (Mannerheim, 1844)	2	0	3	0	1	6
<i>Enicmus fungicola</i> Thomson, C.G., 1868	1	0	10	2	0	13

Yorkshire Coleoptera checklist 2011

<i>Enicmus histrio</i> Joy & Tomlin, 1910	8	8	8	4	2	30
<i>Enicmus rugosus</i> (Herbst, 1793)	0	4	1	1	0	6
<i>Enicmus testaceus</i> (Stephens, 1830)	2	1	6	2	0	11
<i>Enicmus transversus</i> (Olivier, 1790)	68	23	185	33	12	321
<i>Dienerella elongata</i> (Curtis, 1830)	0	1	6	1	0	8
<i>Dienerella filiformis</i> (Gyllenhal, 1827)	0	1	2	0	0	3
<i>Dienerella filum</i> (Aubé, 1850)	1	0	3	0	0	4
<i>Dienerella ruficollis</i> (Marsham, 1802)	7	1	7	1	0	16
<i>Corticaria crenulata</i> (Gyllenhal, 1827)	18	1	7	0	0	26
<i>Corticaria elongata</i> (Gyllenhal, 1827)	10	9	28	5	0	52
<i>Corticaria fagi</i> Wollaston, 1854	0	0	1	0	0	1
<i>Corticaria ferruginea</i> Marsham, 1802	1	1	3	0	0	5
<i>Corticaria fulva</i> (Comolli, 1837)	1	1	4	1	0	7
<i>Corticaria impressa</i> (Olivier, 1790)	67	10	103	12	0	192
<i>Corticaria inconspicua</i> Wollaston, 1860	0	0	2	0	0	2
<i>Corticaria punctulata</i> Marsham, 1802	14	2	29	2	0	47
<i>Corticaria rubripes</i> Mannerheim, 1844	0	1	1	0	0	2
<i>Corticaria serrata</i> (Paykull, 1798)	1	1	0	0	0	2
<i>Corticaria umbilicata</i> (Beck, 1817)	1	4	1	1	0	7
<i>Corticarina fuscata</i> (Gyllenhal, 1827)	70	62	206	51	15	404
<i>Corticarina similata</i> (Gyllenhal, 1827)	0	4	1	1	0	6
<i>Corticinara gibbosa</i> (Herbst, 1793)	94	74	308	38	7	521
<i>Melanophthalma curticolis</i> (Mannerheim, 1844)	3	0	0	0	0	3
<i>Melanophthalma suturalis</i> (Mannerheim, 1844)	1	0	0	0	0	1

Mycetophagidae [13/10]

<i>Pseudotriphyllus suturalis</i> (Fabricius, 1801)	12	23	20	13	4	72
<i>Triphyllus bicolor</i> (Fabricius, 1777)	0	13	3	1	0	17
<i>Litargus connexus</i> (Geoffroy in Fourcroy, 1785)	8	6	20	17	1	52
<i>Mycetophagus atomarius</i> (Fabricius, 1787)	3	3	21	11	3	41
<i>Mycetophagus multipunctatus</i> Fabricius, 1792	4	0	12	2	0	18
<i>Mycetophagus piceus</i> (Fabricius, 1777)	4	5	14	5	1	29
<i>Mycetophagus populi</i> Fabricius, 1798	0	0	0	2	0	2
<i>Mycetophagus quadriguttatus</i> Müller, P.W.J., 1821	0	0	1	0	0	1
<i>Mycetophagus quadripustulatus</i> (Linnaeus, 1761)	8	16	69	23	5	121
<i>Typhaea stercorea</i> (Linnaeus, 1758)	17	11	39	11	2	80

Ciidae [22/16]

<i>Cis bidentatus</i> (Olivier, 1790)	6	10	14	19	1	50
<i>Cis bilamellatus</i> Wood, 1884	19	12	70	26	0	127
<i>Cis boleti</i> (Scopoli, 1763)	9	20	135	35	8	207
<i>Cis fagi</i> Walth, 1839	2	1	0	1	0	4
<i>Cis festivus</i> (Panzer, 1793)	0	5	5	2	1	13
<i>Cis hispidus</i> (Paykull, 1798)	3	0	1	0	0	4
<i>Cis lineatocribratus</i> Mellié, 1849	0	0	2	0	0	2
<i>Cis micans</i> (Fabricius, 1792)	0	0	1	0	0	1
<i>Cis nitidus</i> (Fabricius, 1792)	4	13	26	7	0	50
<i>Cis pygmaeus</i> (Marsham, 1802)	0	1	0	1	0	2
<i>Cis vestitus</i> Mellié, 1849	0	11	2	3	1	17
<i>Cis villosulus</i> (Marsham, 1802)	1	0	3	1	0	5

Yorkshire Coleoptera checklist 2011

<i>Ennearthron cornutum</i> (Gyllenhal, 1827)	2	1	6	1	0	10
<i>Orthocis alni</i> (Gyllenhal, 1813)	1	7	7	3	1	19
<i>Sulcacis affinis</i> (Gyllenhal, 1827)	0	4	4	0	0	8
<i>Octotemnus glabriculus</i> (Gyllenhal, 1827)	4	11	75	27	4	121

Tetratomidae [4/4]

<i>Hallomenus binotatus</i> (Quensel, 1790)	0	3	15	8	1	27
<i>Tetratoma ancora</i> Fabricius, 1791	0	2	0	2	0	4
<i>Tetratoma desmarestii</i> Latreille, 1807	1	4	3	0	0	8
<i>Tetratoma fungorum</i> Fabricius, 1790	9	14	68	16	3	110

Melandryidae [17/11]

<i>Orchesia micans</i> (Panzer, 1793)	4	7	8	6	0	25
<i>Orchesia minor</i> Walker, 1836	0	10	6	0	4	20
<i>Orchesia undulata</i> Kraatz, 1853	1	16	48	7	4	76
<i>Anisoxya fuscata</i> (Illiger, 1798)	0	2	2	0	0	4
<i>Abdera flexuosa</i> (Paykull, 1799)	0	9	1	1	1	12
<i>Abdera quadrifasciata</i> (Curtis, 1829)	0	2	0	0	0	2
<i>Phloiotrya vaudoueri</i> Mulsant, 1856	1	0	4	0	0	5
<i>Hypulus quercinus</i> (Quensel, 1790)	0	3	2	0	0	5
<i>Melandrya caraboides</i> (Linnaeus, 1761)	1	10	9	5	1	26
<i>Conopalpus testaceus</i> (Olivier, 1790)	0	3	6	1	0	10
<i>Osphya bipunctata</i> (Fabricius, 1775)	0	1	0	0	0	1

Mordellidae [17/4]

<i>Mordellistena (Mordellistena) neuwaldeggiana</i> (Panzer, 1796)	0	1	2	1	0	4
<i>Mordellistena (Mordellistena) pumila</i> (Gyllenhal, 1810)	2	0	25	1	0	28
<i>Mordellistena (Mordellistena) variegata</i> (Fabricius, 1798)	0	0	1	0	0	1
<i>Mordellochroa abdominalis</i> (Fabricius, 1775)	0	5	2	0	0	7

Ripiphoridae [1/1]

<i>Metoecus paradoxus</i> (Linnaeus, 1761)	3	9	9	6	2	29
--	---	---	---	---	---	----

Colydiidae [12/5]

<i>Orthocerus clavicornis</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Synchita humeralis</i> (Fabricius, 1792)	1	0	6	1	0	8
<i>Synchita separanda</i> (Reitter, 1882)	0	0	1	0	0	1
<i>Bitoma crenata</i> (Fabricius, 1775)	6	5	27	6	0	44
<i>Colydium elongatum</i> (Fabricius, 1787)	0	0	1	0	0	1

Tenebrionidae [47/31]

<i>Lagria hirta</i> (Linnaeus, 1758)	55	23	123	28	35	264
<i>Eledona agricola</i> (Herbst, 1783)	5	13	14	2	0	34
<i>Tenebrio molitor</i> Linnaeus, 1758	8	6	12	4	1	31
<i>Tenebrio obscurus</i> Fabricius, 1792	0	1	0	0	0	1
<i>Alphitobius diaperinus</i> (Panzer, 1796)	1	1	6	1	1	10
<i>Alphitobius laevigatus</i> (Fabricius, 1781)	1	1	2	0	0	4
<i>Tribolium castaneum</i> (Herbst, 1797)	3	4	8	3	1	19
<i>Tribolium confusum</i> Jacquelin du Val, 1863	0	1	4	0	0	5
<i>Palorus ratzeburgii</i> (Wissmann, 1848)	1	1	0	1	0	3

Yorkshire Coleoptera checklist 2011

<i>Phylan gibbus</i> (Fabricius, 1775)	4	1	0	0	0	5
<i>Melanimon tibialis</i> (Fabricius, 1781)	0	1	6	1	0	8
<i>Nalassus laevioctostriatus</i> (Goeze, 1777)	0	3	10	0	0	13
<i>Xanthomus pallidus</i> (Curtis, 1830)	3	0	0	0	0	3
<i>Blaps mortisaga</i> (Linnaeus, 1758)	0	1	1	0	0	2
<i>Blaps mucronata</i> Latreille, 1804	3	11	8	1	1	24
<i>Prionychus melanarius</i> (Germar, 1813)	0	0	1	0	0	1
<i>Gonodera luperus</i> (Herbst, 1783)	0	0	2	2	0	4
<i>Pseudocistela ceramboides</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Isomira murina</i> (Linnaeus, 1758)	9	7	12	4	2	34
<i>Mycetochara humeralis</i> (Fabricius, 1787)	0	0	1	0	0	1
<i>Crypticus quisquilius</i> (Linnaeus, 1761)	14	0	0	0	0	14
<i>Phaleria cadaverina</i> (Fabricius, 1792)	5	0	0	0	0	5
<i>Myrmechixenus subterraneus</i> Chevrolat, 1835	0	0	1	0	0	1
<i>Myrmechixenus vaporariorum</i> Guérin-Méneville, 1843	0	0	0	1	0	1
<i>Corticeus bicolor</i> (Olivier, 1790)	0	5	9	1	0	15
<i>Corticeus fraxini</i> (Kugelann, 1794)	0	0	1	0	0	1
<i>Corticeus linearis</i> (Fabricius, 1790)	1	0	0	0	0	1
<i>Corticeus unicolor</i> Piller & Mitterpacher, 1783	6	0	15	1	1	23
<i>Scaphidema metallicum</i> (Fabricius, 1792)	25	11	25	3	0	64
<i>Alphitophagus bifasciatus</i> (Say, 1823)	3	1	4	0	0	8
<i>Gnatocerus cornutus</i> (Fabricius, 1798)	2	4	2	0	0	8

Oedemeridae [10/8]

<i>Nacerdes melanura</i> (Linnaeus, 1758)	18	4	7	6	0	35
<i>Ischnomera caerulea</i> (Linnaeus, 1758)	0	4	0	1	0	5
<i>Ischnomera cinerascens</i> (Pandellé, 1867)	0	3	0	0	0	3
<i>Ischnomera cyanea</i> (Fabricius, 1792)	0	8	0	1	0	9
<i>Ischnomera sanguinicollis</i> (Fabricius, 1787)	0	6	0	0	2	8
<i>Oedemera (Oedemera) lurida</i> (Marsham, 1802)	24	3	140	11	7	185
<i>Oedemera (Oedemera) nobilis</i> (Scopoli, 1763)	0	4	6	1	0	11
<i>Oedemera (Oedemera) virescens</i> (Linnaeus, 1767)	0	28	0	0	0	28

Meloidae [10/2]

<i>Meloe proscarabaeus</i> Linnaeus, 1758	1	8	6	5	2	22
<i>Meloe violaceus</i> Marsham, 1802	2	6	20	4	3	35

Pyrochroidae [3/2]

<i>Pyrochroa coccinea</i> (Linnaeus, 1761)	1	10	8	0	0	19
<i>Pyrochroa serraticornis</i> (Scopoli, 1763)	38	18	43	26	11	136

Salpingidae [11/9]

<i>Aglenus brunneus</i> (Gyllenhal, 1813)	0	0	1	0	0	1
<i>Lissodema cursor</i> (Gyllenhal, 1813)	0	0	0	2	0	2
<i>Lissodema denticolle</i> (Gyllenhal, 1813)	0	1	0	0	0	1
<i>Rabocerus foveolatus</i> (Ljungh, 1824)	0	1	2	0	0	3
<i>Rabocerus gabrieli</i> Gerhardt, 1901	2	3	9	6	0	20
<i>Sphaeriestes castaneus</i> (Panzer, 1796)	1	8	6	0	1	16
<i>Sphaeriestes reyi</i> (Abeille, 1874)	5	4	3	1	1	14
<i>Vincenzellus ruficollis</i> (Panzer, 1794)	10	22	41	20	4	97

Yorkshire Coleoptera checklist 2011

Salpingus planirostris (Fabricius, 1787) 19 49 89 19 5 181

Anthicidae [13/7]

Notoxus monoceros (Linnaeus, 1761) 32 3 0 3 0 38
 Anthicus antherinus (Linnaeus, 1761) 2 0 2 0 0 4
 Cyclodinus constrictus (Curtis, 1838) 1 1 0 0 0 2
 Omonadus floralis (Linnaeus, 1758) 39 15 56 18 3 131
 Omonadus formicarius (Goeze, 1777) 13 8 26 7 5 59
 Cordicomus instabilis (Schmidt, 1842) 0 1 0 0 0 1
 Stricticomus tobias (Marseul, 1879) 1 0 5 0 0 6

Aderidae [3/3]

Aderus populneus (Creutzer in Panzer, 1796) 2 0 8 0 0 10
 Eugenes oculus (Paykull, 1798) 1 10 2 6 0 19
 Vanonus brevicornis (Perris, 1869) 0 0 1 0 0 1

Scraptiidae [14/10]

Anaspis (Anaspis) frontalis (Linnaeus, 1758) 10 36 121 24 10 201
 Anaspis (Anaspis) garneysi Fowler, 1889 3 8 7 5 0 23
 Anaspis (Anaspis) humeralis (Fabricius, 1775) 17 25 64 21 4 131
 Anaspis (Anaspis) lurida Stephens, 1832 0 1 2 3 1 7
 Anaspis (Anaspis) maculata (Geoffrey in Fourcroy, 1785) 71 74 352 56 20 573
 Anaspis (Anaspis) pulicaria Costa, A., 1854 0 4 0 2 0 6
 Anaspis (Anaspis) regimbarti Schilsky, 1895 24 24 112 12 4 176
 Anaspis (Anaspis) thoracica (Linnaeus, 1758) 1 2 6 1 0 10
 Anaspis (Nassipa) costai Emery, 1876 2 3 11 0 0 16
 Anaspis (Nassipa) rufilabris (Gyllenhal, 1827) 13 36 121 16 10 196

Cerambycidae [65/47]

Rhagium (Rhagium) inquisitor (Linnaeus, 1758) 2 4 2 0 0 8
 Rhagium (Hagrium) bifasciatum Fabricius, 1775 2 22 49 8 5 86
 Rhagium (Megarhagium) mordax (De Geer, 1775) 1 22 34 16 4 77
 Stenocorus meridianus (Linnaeus, 1758) 5 13 16 12 3 49
 Grammoptera abdominalis (Stephens, 1831) 1 0 3 1 0 5
 Grammoptera ruficornis (Fabricius, 1781) 38 45 263 45 13 404
 Leptura quadrifasciata Linnaeus, 1758 11 19 31 4 0 65
 Anastrangalia sanguinolenta (Linnaeus, 1761) 0 0 1 0 0 1
 Stictoleptura rubra (Linnaeus, 1758) 1 0 2 0 0 3
 Stictoleptura scutellata (Fabricius, 1781) 0 0 2 0 0 2
 Anoplodera sexguttata (Fabricius, 1775) 0 9 0 0 0 9
 Pachytodes cerambyciformis (Schränk, 1781) 0 0 4 0 0 4
 Alosterna tabacicolor (De Geer, 1775) 0 27 6 2 0 35
 Rutpela maculata (Poda, 1761) 20 14 111 11 0 156
 Stenurella melanura (Linnaeus, 1758) 0 6 2 0 0 8
 Asemum striatum (Linnaeus, 1758) 1 3 2 2 0 8
 Tetropium gabrieli Weise, 1905 0 4 4 2 0 10
 Arhopalus fesus (Mulsant, 1839) 1 0 0 0 0 1
 Arhopalus rusticus (Linnaeus, 1758) 2 0 1 0 0 3
 Cerambyx cerdo Linnaeus, 1758 1 0 0 0 0 1
 Cerambyx scopolii Fuessly, 1775 0 1 2 0 0 3

Yorkshire Coleoptera checklist 2011

Gracilia minuta (Fabricius, 1781)	2	0	4	2	0	8
Nathrius brevipennis (Mulsant, 1839)	0	0	3	1	0	4
Molorchus minor (Linnaeus, 1758)	1	1	0	3	0	5
Glaphyra umbellatarum (von Schreber, 1759)	0	0	0	2	0	2
Aromia moschata (Linnaeus, 1758)	1	2	2	3	0	8
Hylotrupes bajulus (Linnaeus, 1758)	0	0	2	0	0	2
Callidium violaceum (Linnaeus, 1758)	1	0	1	0	0	2
Pyrrhidium sanguineum (Linnaeus, 1758)	0	0	0	1	0	1
Phymatodes testaceus (Linnaeus, 1758)	2	10	1	2	0	15
Poecilium alni (Linnaeus, 1767)	0	0	2	0	0	2
Clytus arietis (Linnaeus, 1758)	5	27	96	11	1	140
Plagionotus arcuatus (Linnaeus, 1758)	0	0	1	1	0	2
Anaglyptus mysticus (Linnaeus, 1758)	6	2	13	3	0	24
Mesosa nebulosa (Fabricius, 1781)	0	0	2	0	0	2
Agapanthia villosviridescens (De Geer, 1775)	19	0	36	0	1	56
Pogonocherus fasciculatus (De Geer, 1775)	0	1	0	1	0	2
Pogonocherus hispidulus (Piller & Mitterpacher, 1783)	4	18	5	3	3	33
Pogonocherus hispidus (Linnaeus, 1758)	1	10	1	3	1	16
Acanthocinus aedilis (Linnaeus, 1758)	2	1	6	1	0	10
Leiopus nebulosus (Linnaeus, 1758)	5	22	32	19	0	78
Saperda carcharias (Linnaeus, 1758)	0	3	7	0	0	10
Saperda populnea (Linnaeus, 1758)	5	6	15	11	0	37
Saperda scalaris (Linnaeus, 1758)	0	8	5	0	0	13
Stenostola dubia (Laicharting, 1784)	0	6	7	9	3	25
Phytoecia cylindrica (Linnaeus, 1758)	2	0	0	1	0	3
Tetrops praeustus (Linnaeus, 1758)	2	6	22	6	0	36

Megalopodidae [3/1]

Zeugophora subspinosa (Fabricius, 1781)	4	3	16	1	0	24
---	---	---	----	---	---	----

Orsodacnidae [2/1]

Orsodacne cerasi (Linnaeus, 1758)	0	31	7	8	0	46
-----------------------------------	---	----	---	---	---	----

Chrysomelidae [270/211]

Bruchus loti Paykull, 1800	37	24	134	15	0	210
Bruchus pisorum (Linnaeus, 1758)	2	1	2	0	0	5
Bruchus rufimanus Boheman, 1833	15	8	30	1	0	54
Bruchus rufipes Herbst, 1783	0	0	4	0	0	4
Bruchidius cisti (Fabricius, 1775)	5	11	9	3	0	28
Bruchidius varius (Olivier, 1795)	3	0	2	0	0	5
Bruchidius villosus (Fabricius, 1792)	4	2	63	9	0	78
Callosobruchus chinensis (Linnaeus, 1758)	0	1	0	0	0	1
Macrolea appendiculata (Panzer, 1794)	4	1	0	9	0	14
Macrolea mutica (Fabricius, 1792)	4	0	4	0	0	8
Donacia aquatica (Linnaeus, 1758)	0	1	1	1	0	3
Donacia bicolora Zschach, 1788	0	2	0	0	0	2
Donacia cinerea Herbst, 1784	7	3	6	1	0	17
Donacia clavipes Fabricius, 1792	12	2	3	3	0	20
Donacia crassipes Fabricius, 1775	0	0	1	0	0	1
Donacia dentata Hoppe, 1795	1	0	0	0	0	1

Yorkshire Coleoptera checklist 2011

<i>Donacia impressa</i> Paykull, 1799	5	1	1	0	0	7
<i>Donacia marginata</i> Hoppe, 1795	1	1	1	5	0	8
<i>Donacia semicuprea</i> Panzer, 1796	12	2	21	1	0	36
<i>Donacia simplex</i> Fabricius, 1775	18	11	36	13	2	80
<i>Donacia sparganii</i> Ahrens, 1810	7	0	0	2	0	9
<i>Donacia thalassina</i> Germar, 1811	4	6	4	0	0	14
<i>Donacia versicolore</i> a (Brahm, 1791)	14	5	15	11	0	45
<i>Donacia vulgaris</i> Zschach, 1788	14	9	20	6	0	49
<i>Plateumaris affinis</i> (Kunze, 1818)	0	3	2	2	4	11
<i>Plateumaris braccata</i> (Scopoli, 1772)	4	0	0	0	0	4
<i>Plateumaris discolor</i> (Panzer, 1795)	0	21	21	14	5	61
<i>Plateumaris sericea</i> (Linnaeus, 1758)	16	12	17	8	1	54
<i>Lema cyanella</i> (Linnaeus, 1758)	4	6	2	2	0	14
<i>Oulema melanopus</i> (Linnaeus, 1758)	6	0	5	4	1	16
<i>Oulema obscura</i> (Stephens, 1831)	13	30	11	17	1	72
<i>Oulema rufocyanea</i> (Suffrian, 1847)	37	11	48	14	6	116
<i>Crioceris asparagi</i> (Linnaeus, 1758)	0	0	20	0	0	20
<i>Lilioceris lili</i> (Scopoli, 1763)	7	0	9	3	0	19
<i>Labidostomis tridentata</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Clytra quadripunctata</i> (Linnaeus, 1758)	0	12	16	0	0	28
<i>Cryptocephalus aureolus</i> Suffrian, 1847	2	11	0	0	0	13
<i>Cryptocephalus biguttatus</i> (Scopoli, 1763)	0	2	0	0	0	2
<i>Cryptocephalus bipunctatus</i> (Linnaeus, 1758)	0	1	0	1	0	2
<i>Cryptocephalus fulvus</i> (Goeze, 1777)	2	2	20	2	0	26
<i>Cryptocephalus hypochaeridis</i> (Linnaeus, 1758)	0	2	0	0	0	2
<i>Cryptocephalus labiatus</i> (Linnaeus, 1761)	4	13	85	12	0	114
<i>Cryptocephalus moraei</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Cryptocephalus parvulus</i> Müller, O.F., 1776	0	0	8	0	0	8
<i>Cryptocephalus punctiger</i> Paykull, 1799	0	0	2	0	0	2
<i>Cryptocephalus pusillus</i> Fabricius, 1777	13	10	34	6	0	63
<i>Oomorplus concolor</i> (Sturm, 1807)	0	1	0	1	0	2
<i>Timarcha goettingensis</i> (Linnaeus, 1758)	2	1	27	8	1	39
<i>Timarcha tenebricosa</i> (Fabricius, 1775)	0	0	9	3	0	12
<i>Leptinotarsa decemlineata</i> (Say, 1824)	0	0	1	0	0	1
<i>Chrysolina americana</i> (Linnaeus, 1758)	2	0	0	0	0	2
<i>Chrysolina banksi</i> (Fabricius, 1775)	0	0	2	0	0	2
<i>Chrysolina brunsvicensis</i> (Gravenhorst, 1807)	0	5	0	3	2	10
<i>Chrysolina fastuosa</i> (Scopoli, 1763)	0	0	28	0	0	28
<i>Chrysolina graminis</i> (Linnaeus, 1758)	5	12	1	15	0	33
<i>Chrysolina herbacea</i> (Duftschmid, 1825)	0	0	1	1	0	2
<i>Chrysolina hyperici</i> (Forster, 1771)	2	6	23	2	0	33
<i>Chrysolina marginata</i> (Linnaeus, 1758)	1	1	3	0	2	7
<i>Chrysolina oricalcia</i> (Müller, O.F., 1776)	3	1	10	5	5	24
<i>Chrysolina polita</i> (Linnaeus, 1758)	29	27	96	26	5	183
<i>Chrysolina sanguinolenta</i> (Linnaeus, 1758)	1	0	7	0	0	8
<i>Chrysolina staphylaea</i> (Linnaeus, 1758)	10	13	29	11	4	67
<i>Chrysolina varians</i> (Schaller, 1783)	2	13	3	4	2	24
<i>Chrysolina violacea</i> (Müller, O.F., 1776)	0	1	3	2	0	6
<i>Gastrophysa polygoni</i> (Linnaeus, 1758)	20	10	50	31	2	113
<i>Gastrophysa viridula</i> (De Geer, 1775)	28	20	143	32	5	228

Yorkshire Coleoptera checklist 2011

<i>Phaedon armoraciae</i> (Linnaeus, 1758)	21	22	56	16	3	118
<i>Phaedon cochleariae</i> (Fabricius, 1792)	27	12	43	18	10	110
<i>Phaedon concinnus</i> Stephens, 1831	10	0	0	0	0	10
<i>Phaedon tumidulus</i> (Germar, 1824)	53	32	177	26	4	292
<i>Hydrothassa glabra</i> (Herbst, 1783)	3	5	3	9	1	21
<i>Hydrothassa hannoveriana</i> (Fabricius, 1775)	0	0	0	9	2	11
<i>Hydrothassa marginella</i> (Linnaeus, 1758)	15	35	42	29	5	126
<i>Prasocuris junci</i> (Brahm, 1790)	11	9	6	5	1	32
<i>Prasocuris phellandrii</i> (Linnaeus, 1758)	5	2	2	3	1	13
<i>Plagioderia versicolora</i> (Laicharting, 1781)	2	3	4	0	0	9
<i>Chrysomela aenea</i> Linnaeus, 1758	1	39	9	9	5	63
<i>Chrysomela populi</i> Linnaeus, 1758	10	10	7	1	0	28
<i>Gonioctena decemnotata</i> (Marsham, 1802)	0	0	1	0	0	1
<i>Gonioctena olivacea</i> (Forster, 1771)	0	4	7	0	1	12
<i>Gonioctena pallida</i> (Linnaeus, 1758)	0	33	2	9	4	48
<i>Gonioctena viminalis</i> (Linnaeus, 1758)	0	3	0	0	0	3
<i>Phratora laticollis</i> Suffrian, 1851	12	6	11	3	0	32
<i>Phratora vitellinae</i> (Linnaeus, 1758)	14	5	31	29	12	91
<i>Phratora vulgatissima</i> (Linnaeus, 1758)	51	13	110	23	18	215
<i>Galerucella calmariensis</i> (Linnaeus, 1767)	9	3	20	5	0	37
<i>Galerucella lineola</i> (Fabricius, 1781)	9	12	73	15	7	116
<i>Galerucella nymphaeae</i> (Linnaeus, 1758)	7	6	70	8	0	91
<i>Galerucella pusilla</i> (Duftschmid, 1825)	2	0	9	6	0	17
<i>Galerucella sagittariae</i> (Gyllenhal, 1813)	27	6	17	8	0	58
<i>Galerucella tenella</i> (Linnaeus, 1761)	14	17	16	29	2	78
<i>Pyrrhalta viburni</i> (Paykull, 1799)	1	6	23	4	0	34
<i>Xanthogaleruca luteola</i> (Müller, O.F., 1766)	0	0	1	0	0	1
<i>Galeruca tanacetii</i> (Linnaeus, 1758)	8	3	16	3	0	30
<i>Lochmaea caprea</i> (Linnaeus, 1758)	7	9	100	14	1	131
<i>Lochmaea crataegi</i> (Forster, 1771)	6	19	79	14	5	123
<i>Lochmaea suturalis</i> (Thomson, C.G., 1866)	9	27	129	7	6	178
<i>Phyllotreta quadrimaculata</i> (Linnaeus, 1758)	2	7	12	12	0	33
<i>Luperus flavipes</i> (Linnaeus, 1767)	1	6	15	0	0	22
<i>Luperus longicornis</i> (Fabricius, 1781)	8	21	13	16	5	63
<i>Calomicrus circumfusus</i> (Marsham, 1802)	0	0	1	0	0	1
<i>Agelastica alni</i> (Linnaeus, 1758)	0	0	1	0	0	1
<i>Sermylassa halensis</i> (Linnaeus, 1767)	20	12	29	2	2	65
<i>Phyllotreta atra</i> (Fabricius, 1775)	3	9	6	0	0	18
<i>Phyllotreta consobrina</i> (Curtis, 1837)	0	2	0	0	0	2
<i>Phyllotreta cruciferae</i> (Goeze, 1777)	0	3	0	0	0	3
<i>Phyllotreta diademata</i> Foudras, 1860	0	0	5	0	0	5
<i>Phyllotreta exclamationis</i> (Thunberg, 1784)	9	3	7	4	1	24
<i>Phyllotreta flexuosa</i> (Illiger, 1794)	2	3	2	10	2	19
<i>Phyllotreta nemorum</i> (Linnaeus, 1758)	4	2	10	14	7	37
<i>Phyllotreta nigripes</i> (Fabricius, 1775)	8	6	42	6	0	62
<i>Phyllotreta nodicornis</i> (Marsham, 1802)	4	1	19	1	1	26
<i>Phyllotreta ochripes</i> (Curtis, 1837)	8	3	3	0	0	14
<i>Phyllotreta punctulata</i> (Marsham, 1802)	1	1	1	0	0	3
<i>Phyllotreta tetrastigma</i> (Comolli, 1837)	12	6	26	15	1	60
<i>Phyllotreta undulata</i> Kutschera, 1860	29	39	78	26	4	176

Yorkshire Coleoptera checklist 2011

<i>Phyllotreta vittula</i> (Redtenbacher, 1849)	1	3	7	3	0	14
<i>Aphthona ?atratula</i> Allard, 1859	1	1	9	1	0	12
<i>Aphthona atrocaerulea</i> (Stephens, 1829)	5	3	15	1	0	24
<i>Aphthona euphorbiae</i> (Schrank, 1781)	71	43	264	25	4	407
<i>Aphthona herbigrada</i> (Curtis, 1837)	3	7	0	5	1	16
<i>Aphthona lutescens</i> (Gyllenhal, 1808)	0	0	2	0	0	2
<i>Aphthona melancholica</i> Weise, 1888	1	1	8	2	0	12
<i>Aphthona nigriceps</i> (Redtenbacher, 1842)	1	2	1	10	0	14
<i>Aphthona nonstriata</i> (Goeze, 1777)	9	7	21	32	1	70
<i>Longitarsus aeneicollis</i> (Faldermann, 1837)	2	2	3	1	0	8
<i>Longitarsus anchusae</i> (Paykull, 1799)	0	2	5	0	0	7
<i>Longitarsus atricillus</i> (Linnaeus, 1761)	3	5	5	0	1	14
<i>Longitarsus brunneus</i> (Duftschmid, 1825)	0	2	2	3	0	7
<i>Longitarsus dorsalis</i> (Fabricius, 1781)	0	1	0	0	0	1
<i>Longitarsus exoletus</i> (Linnaeus, 1758)	3	1	11	1	0	16
<i>Longitarsus flavicornis</i> (Stephens, 1831)	0	0	3	0	0	3
<i>Longitarsus ganglbaueri</i> Heikertinger, 1912	1	3	2	0	2	8
<i>Longitarsus gracilis</i> Kutschera, 1864	21	9	10	5	2	47
<i>Longitarsus holsaticus</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Longitarsus jacobaeae</i> (Waterhouse, G.R., 1858)	26	8	63	10	6	113
<i>Longitarsus kutscherae</i> (Rye, 1872)	4	2	16	0	0	22
<i>Longitarsus luridus</i> (Scopoli, 1763)	59	34	199	43	7	342
<i>Longitarsus melanocephalus</i> (De Geer, 1775)	11	5	44	8	0	68
<i>Longitarsus membranaceus</i> (Foudras, 1860)	2	5	0	0	4	11
<i>Longitarsus nasturtii</i> (Fabricius, 1792)	0	1	0	1	0	2
<i>Longitarsus nigerrimus</i> (Gyllenhal, 1827)	0	1	0	0	0	1
<i>Longitarsus nigrofasciatus</i> (Goeze, 1777)	1	0	1	0	0	2
<i>Longitarsus obliteratus</i> (Rosenhauer, 1847)	2	0	1	0	0	3
<i>Longitarsus ochroleucus</i> (Marsham, 1802)	3	2	7	2	0	14
<i>Longitarsus parvulus</i> (Paykull, 1799)	22	18	49	5	0	94
<i>Longitarsus pellucidus</i> (Foudras, 1860)	2	1	7	0	0	10
<i>Longitarsus pratensis</i> (Panzer, 1794)	20	5	9	4	0	38
<i>Longitarsus reichei</i> (Allard, 1860)	3	0	11	0	0	14
<i>Longitarsus rubiginosus</i> (Foudras, 1860)	6	0	17	0	0	23
<i>Longitarsus succineus</i> (Foudras, 1860)	15	7	41	5	3	71
<i>Longitarsus suturellus</i> (Duftschmid, 1825)	25	14	89	22	2	152
<i>Longitarsus tabidus</i> (Fabricius, 1775)	0	1	0	0	0	1
<i>Altica brevicollis</i> Foudras, 1860	0	1	1	0	0	2
<i>Altica ericeti</i> (Allard, 1859)	1	10	13	2	3	29
<i>Altica helianthemi</i> (Allard, 1859)	3	5	10	4	0	22
<i>Altica lythri</i> Aubé, 1843	58	44	262	34	16	414
<i>Altica oleracea</i> (Linnaeus, 1758)	12	10	45	7	0	74
<i>Altica palustris</i> Weise, 1888	16	11	69	13	2	111
<i>Hermaeophaga mercurialis</i> (Fabricius, 1792)	0	0	1	0	0	1
<i>Batophila aerata</i> (Marsham, 1802)	0	0	0	1	0	1
<i>Batophila rubi</i> (Paykull, 1799)	3	23	43	11	3	83
<i>Lythriaria salicariae</i> (Paykull, 1800)	0	1	2	0	0	3
<i>Ochrosis ventralis</i> (Illiger, 1807)	0	0	1	0	0	1
<i>Neocrepidodera ferruginea</i> (Scopoli, 1763)	63	29	190	36	10	328
<i>Neocrepidodera impressa</i> (Fabricius, 1801)	1	0	0	0	0	1

Yorkshire Coleoptera checklist 2011

<i>Neocrepidodera transversa</i> (Marsham, 1802)	15	23	121	21	5	185
<i>Derocrepis rufipes</i> (Linnaeus, 1758)	9	17	23	15	6	70
<i>Hippuriphila modeeri</i> (Linnaeus, 1761)	13	12	14	5	4	48
<i>Crepidodera aurata</i> (Marsham, 1802)	14	4	11	4	0	33
<i>Crepidodera aurea</i> (Fourcroy, 1785)	0	2	12	1	1	16
<i>Crepidodera fulvicornis</i> (Fabricius, 1792)	86	40	320	52	23	521
<i>Crepidodera plutus</i> (Latreille, 1804)	2	0	9	0	0	11
<i>Epitrix atropae</i> Foudras, 1860	2	0	1	0	0	3
<i>Epitrix pubescens</i> (Koch, J.D.W., 1803)	3	0	13	2	0	18
<i>Podagrica fuscicornis</i> (Linnaeus, 1767)	0	3	0	0	0	3
<i>Mantura matthewsii</i> (Curtis, 1833)	4	5	0	8	0	17
<i>Mantura obtusata</i> (Gyllenhal, 1813)	0	4	1	7	0	12
<i>Mantura rustica</i> (Linnaeus, 1767)	5	6	13	8	0	32
<i>Chaetocnema arida</i> Foudras, 1860	1	2	0	0	0	3
<i>Chaetocnema concinna</i> (Marsham, 1802)	49	29	216	46	8	348
<i>Chaetocnema confusa</i> (Boheman, 1851)	3	0	1	0	0	4
<i>Chaetocnema hortensis</i> (Fourcroy, 1785)	15	11	45	19	8	98
<i>Chaetocnema sahlbergii</i> (Gyllenhal, 1827)	0	1	0	0	0	1
<i>Chaetocnema subcoerulea</i> (Kutschera, 1864)	0	1	0	0	0	1
<i>Sphaeroderma rubidum</i> (Graëlls, 1858)	16	12	27	2	1	58
<i>Sphaeroderma testaceum</i> (Fabricius, 1775)	28	16	79	19	5	147
<i>Apteropeda globosa</i> (Illiger, 1794)	0	3	1	2	0	6
<i>Apteropeda orbiculata</i> (Marsham, 1802)	2	7	7	1	1	18
<i>Mniophila muscorum</i> (Koch, J.D.W., 1803)	0	3	0	3	1	7
<i>Psylliodes affinis</i> (Paykull, 1799)	32	4	111	23	1	171
<i>Psylliodes attenuata</i> (Koch, J.D.W., 1803)	0	0	3	1	0	4
<i>Psylliodes chalcomera</i> (Illiger, 1807)	0	0	1	0	0	1
<i>Psylliodes chrysocephala</i> (Linnaeus, 1758)	20	15	149	7	8	199
<i>Psylliodes cuprea</i> (Koch, J.D.W., 1803)	3	10	12	1	1	27
<i>Psylliodes dulcamarae</i> (Koch, J.D.W., 1803)	1	0	5	0	0	6
<i>Psylliodes hyoscyami</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Psylliodes laticollis</i> Kutschera, 1864	11	4	15	0	0	30
<i>Psylliodes marcida</i> (Illiger, 1807)	7	3	0	0	0	10
<i>Psylliodes napi</i> (Fabricius, 1792)	8	10	26	13	6	63
<i>Psylliodes picina</i> (Marsham, 1802)	15	6	29	9	0	59
<i>Cassida flaveola</i> Thunberg, 1794	4	12	27	14	1	58
<i>Cassida hemisphaerica</i> Herbst, 1799	1	1	0	0	0	2
<i>Cassida nobilis</i> Linnaeus, 1758	0	1	2	0	0	3
<i>Cassida prasina</i> Illiger, 1798	0	0	1	0	0	1
<i>Cassida rubiginosa</i> Müller, O.F., 1776	44	22	155	31	13	265
<i>Cassida vibex</i> Linnaeus, 1767	1	1	0	0	0	2
<i>Cassida viridis</i> Linnaeus, 1758	5	4	12	3	0	24
<u>Nemonychidae [1/1]</u>						
<i>Cimberis attelaboides</i> (Fabricius, 1787)	0	9	0	1	0	10

Anthribidae [9/7]

<i>Anthribus fasciatus</i> (Forster, 1771)	0	0	3	0	0	3
<i>Anthribus nebulosus</i> (Forster, 1771)	4	14	7	10	2	37
<i>Platyrhinus resinosus</i> (Scopoli, 1763)	5	1	22	2	0	30
<i>Platystomos albinus</i> (Linnaeus, 1758)	0	0	0	1	0	1
<i>Choragus sheppardi</i> Kirby, 1819	3	1	1	1	0	6
<i>Araecerus fasciculatus</i> (De Geer, 1775)	0	0	0	1	0	1
<i>Bruchela rufipes</i> (Olivier, 1790)	2	0	15	0	0	17

Rhynchitidae [18/14]

<i>Involvlus cupreus</i> (Linnaeus, 1758)	0	21	1	0	0	22
<i>Lasiorrhynchites</i> (<i>Lasiorrhynchites</i>) <i>cavifrons</i> (Gyllenhal, 1833)	0	2	4	2	0	8
<i>Lasiorrhynchites</i> (<i>Lasiorrhynchites</i>) <i>olivaceus</i> (Gyllenhal, 1833)	0	1	0	0	0	1
<i>Neocoenorrhinus aeneovirens</i> (Marsham, 1802)	0	12	3	0	0	15
<i>Neocoenorrhinus aequatus</i> (Linnaeus, 1767)	5	5	37	1	0	48
<i>Neocoenorrhinus germanicus</i> (Herbst, 1797)	11	8	21	3	0	43
<i>Neocoenorrhinus interpunctatus</i> (Stephens, 1831)	0	0	3	0	0	3
<i>Neocoenorrhinus pauxillus</i> (Germar, 1824)	0	0	0	1	0	1
<i>Temnocerus longiceps</i> (Thomson, C.G., 1888)	2	4	13	6	0	25
<i>Temnocerus nanus</i> (Paykull, 1792)	13	10	49	12	0	84
<i>Temnocerus tomentosus</i> (Gyllenhal, 1839)	1	1	5	1	0	8
<i>Byctiscus betulae</i> (Linnaeus, 1758)	0	2	0	0	0	2
<i>Deporaus</i> (<i>Deporaus</i>) <i>betulae</i> (Linnaeus, 1758)	9	20	67	10	0	106
<i>Deporaus</i> (<i>Caenorhinus</i>) <i>mannerheimii</i> (Hummel, 1823)	8	15	10	2	0	35

Attelabidae [2/2]

<i>Attelabus nitens</i> (Scopoli, 1763)	9	14	50	6	0	79
<i>Apoderus coryli</i> (Linnaeus, 1758)	0	3	1	1	0	5

Apionidae [87/68]

<i>Apion cruentatum</i> Walton, 1844	7	5	10	3	1	26
<i>Apion frumentarium</i> (Linnaeus, 1758)	41	10	88	15	6	160
<i>Apion haematodes</i> Kirby, 1808	14	6	33	6	3	62
<i>Apion rubens</i> Stephens, 1839	2	3	1	2	1	9
<i>Apion rubiginosum</i> Grill, 1893	0	0	1	0	0	1
<i>Perapion</i> (<i>Perapion</i>) <i>affine</i> (Kirby, 1808)	2	3	3	1	0	9
<i>Perapion</i> (<i>Perapion</i>) <i>curtirostre</i> (Germar, 1817)	28	22	103	20	6	179
<i>Perapion</i> (<i>Perapion</i>) <i>hydrolapathi</i> (Marsham, 1802)	19	14	33	9	0	75
<i>Perapion</i> (<i>Perapion</i>) <i>marchicum</i> (Herbst, 1797)	10	5	25	2	0	42
<i>Perapion</i> (<i>Perapion</i>) <i>violaceum</i> (Kirby, 1808)	30	18	115	28	18	209
<i>Aspidapion</i> (<i>Aspidapion</i>) <i>radiolus</i> (Marsham, 1802)	16	9	20	2	2	49
<i>Aspidapion</i> (<i>Koestlinia</i>) <i>aeneum</i> (Fabricius, 1775)	7	6	18	0	1	32
<i>Ceratapion</i> (<i>Ceratapion</i>) <i>carduorum</i> (Kirby, 1808)	9	7	26	6	0	48
<i>Ceratapion</i> (<i>Ceratapion</i>) <i>gibbirostre</i> (Gyllenhal, 1813)	31	1	47	9	5	93
<i>Ceratapion</i> (<i>Acanephodus</i>) <i>onopordi</i> (Kirby, 1808)	19	3	8	1	2	33
<i>Diplapion confluens</i> (Kirby, 1808)	7	0	1	0	0	8
<i>Diplapion stolidum</i> (Germar, 1817)	1	0	0	0	0	1
<i>Omphalapion hookerorum</i> (Kirby, 1808)	13	2	13	3	1	32
<i>Omphalapion laevigatum</i> (Paykull, 1792)	0	0	0	1	0	1
<i>Exapion</i> (<i>Exapion</i>) <i>fuscirostre</i> (Fabricius, 1775)	1	0	8	0	0	9

Yorkshire Coleoptera checklist 2011

Exapion (Ulapion) ulicis (Forster, 1771)	54	14	77	9	9	163
Kalcapion pallipes (Kirby, 1808)	5	15	5	10	0	35
Melanapion minimum (Herbst, 1797)	0	4	1	0	0	5
Squamapion atomarium (Kirby, W., 1808)	3	1	0	0	0	4
Squamapion vicinum (Kirby, W., 1808)	0	0	1	1	0	2
Malvapion malvae (Fabricius, 1775)	15	0	31	2	0	48
Pseudapion rufirostre (Fabricius, 1775)	4	0	9	0	1	14
Rhopalapion longirostre (Olivier, 1807)	0	0	0	1	0	1
Cyanapion (Cyanapion) spencii (Kirby, 1808)	1	3	10	5	1	20
Cyanapion (Bothryorrhynchapion) afer (Gyllenhal, 1833)	3	2	8	3	1	17
Cyanapion (Bothryorrhynchapion) gyllenhalii (Kirby, 1808)	1	2	1	1	0	5
Eutrichapion (Eutrichapion) ervi (Kirby, 1808)	30	19	76	18	5	148
Eutrichapion (Eutrichapion) viciae (Paykull, 1800)	12	7	61	14	2	96
Eutrichapion (Cnemapion) vorax (Herbst, 1797)	9	5	19	5	0	38
Eutrichapion (Psilocalymma) punctigerum (Paykull, 1792)	1	6	4	1	2	14
Hemitrichapion (Tinocyba) reflexum (Gyllenhal, 1833)	0	1	1	0	0	2
Holotrichapion (Holotrichapion) ononis (Kirby, 1808)	20	4	2	3	1	30
Holotrichapion (Apiops) pisi (Fabricius, 1801)	10	6	15	6	2	39
Holotrichapion (Legaricapion) aethiops (Herbst, 1797)	5	3	4	2	2	16
Oxystoma cerdo (Gerstaecker, 1854)	6	0	15	3	0	24
Oxystoma craccae (Linnaeus, 1767)	1	0	1	1	0	3
Oxystoma pomonae (Fabricius, 1798)	0	0	1	0	0	1
Oxystoma subulatum (Kirby, 1808)	21	31	58	16	2	128
Pirapion immune (Kirby, W., 1808)	2	3	6	0	0	11
Catapion curtisi (Stephens, 1831)	1	0	0	0	0	1
Catapion pubescens (Kirby, W., 1811)	2	1	5	1	1	10
Catapion seniculus (Kirby, W., 1808)	15	6	3	5	4	33
Ischnopterapion (Ischnopterapion) loti (Kirby, 1808)	39	29	88	32	4	192
Ischnopterapion (Ischnopterapion) modestum (Germar, 1817)	0	0	3	1	0	4
Ischnopterapion (Chlorapion) virens (Herbst, 1797)	25	12	19	8	2	66
Protopirapion atratum (Germar, 1817)	2	5	6	0	3	16
Stenopterapion (Stenopterapion) meliloti (Kirby, 1808)	10	3	7	0	0	20
Stenopterapion (Stenopterapion) tenue (Kirby, 1808)	13	2	10	4	0	29
Stenopterapion (Cobosiotherium) scutellare (Kirby, 1811)	0	1	4	0	0	5
Synapion ebeninum (Kirby, 1808)	0	0	0	1	0	1
Betulapion simile (Kirby, 1811)	21	4	66	10	0	101
Protapion apricans (Herbst, 1797)	39	21	154	28	15	257
Protapion assimile (Kirby, 1808)	18	36	43	5	1	103
Protapion difforme (Germar, 1818)	0	0	1	0	0	1
Protapion dissimile (Germar, 1817)	0	0	2	0	0	2
Protapion filirostre (Kirby, 1808)	0	0	3	0	0	3
Protapion fulvipes (Geoffroy in Fourcroy, 1785)	67	51	163	35	4	320
Protapion nigritarse (Kirby, 1808)	21	13	61	13	4	112
Protapion ononidis (Gyllenhal, 1827)	6	2	6	4	0	18
Protapion schoenherri (Boheman, 1839)	0	1	0	1	0	2
Protapion trifolii (Linnaeus, 1768)	13	3	14	1	2	33
Protapion varipes (Germar, 1817)	0	0	2	0	0	2
Pseudoprotapion astragali (Paykull, 1800)	0	0	0	1	0	1

Nanophyidae [2/1]

Nanophyes marmoratus (Goeze, 1777)	11	4	33	7	0	55
------------------------------------	----	---	----	---	---	----

Dryophthoridae [4/4]

Dryophthorus corticalis (Paykull, 1792)	0	0	2	0	0	2
Sitophilus granarius (Linnaeus, 1758)	3	4	5	1	0	13
Sitophilus oryzae (Linnaeus, 1763)	2	5	4	2	0	13
Sitophilus zeamais Motschulsky, 1855	0	0	1	0	0	1

Eirrhinidae [12/9]

Grypus equiseti (Fabricius, 1775)	11	22	23	16	4	76
Notaris acridulus (Linnaeus, 1758)	31	11	76	26	5	149
Notaris aethiops (Fabricius, 1792)	0	1	0	1	0	2
Notaris scirpi (Fabricius, 1792)	3	0	25	5	0	33
Thryogenes festucae (Herbst, 1795)	4	0	0	0	0	4
Thryogenes nereis (Paykull, 1800)	0	1	0	1	0	2
Thryogenes scirrhosus (Gyllenhal, 1836)	3	0	0	0	0	3
Tournotaris bimaculatus (Fabricius, 1787)	16	2	10	0	0	28
Stenopelmus rufinasus Gyllenhal, 1836	4	0	6	1	0	11

Curculionidae [475/312]

Archarius pyrrhoceras (Marsham, 1802)	15	22	90	12	3	142
Archarius salicivorus (Paykull, 1792)	23	13	83	21	11	151
Archarius villosus (Fabricius, 1781)	1	2	16	1	0	20
Curculio betulae (Stephens, 1831)	3	1	24	0	1	29
Curculio glandium Marsham, 1802	11	1	40	0	0	52
Curculio nucum Linnaeus, 1758	0	0	7	0	0	7
Curculio rubidus (Gyllenhal, 1836)	0	1	6	0	0	7
Curculio venosus (Gravenhorst, 1807)	1	1	33	2	0	37
Anthonomus (Anthonomus) bituberculatus Thomson, C.G., 1868	4	3	10	4	3	24
Anthonomus (Anthonomus) conspersus Desbrochers, 1868	0	5	1	0	0	6
Anthonomus (Anthonomus) pedicularius (Linnaeus, 1758)	12	21	54	11	2	100
Anthonomus (Anthonomus) piri Kollar, 1837	0	1	1	0	0	2
Anthonomus (Anthonomus) pomorum (Linnaeus, 1758)	4	5	4	2	0	15
Anthonomus (Anthonomus) rubi (Herbst, 1795)	16	11	44	8	4	83
Anthonomus (Anthonomus) ulmi (De Geer, 1775)	0	3	4	1	4	12
Anthonomus (Furcipes) rectirostris (Linnaeus, 1758)	0	4	0	0	0	4
Cionus alauda (Herbst, 1784)	8	9	23	7	1	48
Cionus hortulanus (Fourcroy, 1785)	0	0	1	0	0	1
Cionus scrophulariae (Linnaeus, 1758)	16	15	90	20	8	149
Cleopus pulchellus (Herbst, 1795)	1	7	7	6	0	21
Elleucus bipunctatus (Linnaeus, 1758)	0	3	0	0	0	3
Dorytomus dejeani Faust, 1882	5	1	28	4	0	38
Dorytomus filirostris (Gyllenhal, 1836)	7	0	5	2	0	14
Dorytomus hirtipennis Bedel, 1884	2	0	1	0	0	3
Dorytomus longimanus (Forster, 1771)	9	5	18	3	0	35
Dorytomus melanophthalmus (Paykull, 1792)	7	7	16	7	0	37
Dorytomus rufatus Bedel, 1888	11	11	32	4	0	58
Dorytomus salicinus (Gyllenhal, 1827)	1	0	2	7	0	10
Dorytomus salicis Walton, 1851	3	1	2	0	0	6

Yorkshire Coleoptera checklist 2011

<i>Dorytomus taeniatus</i> (Fabricius, 1781)	11	13	75	15	10	124
<i>Dorytomus tortrix</i> (Linnaeus, 1761)	6	2	24	3	1	36
<i>Dorytomus tremulae</i> (Fabricius, 1787)	1	0	9	1	0	11
<i>Cleopomiarus graminis</i> (Gyllenhal, 1813)	1	0	3	1	0	5
<i>Cleopomiarus plantarum</i> (Germar, 1824)	0	0	1	0	0	1
<i>Gymnetron beccabungae</i> (Linnaeus, 1761)	0	0	1	0	0	1
<i>Gymnetron melanarium</i> (Germar, 1821)	3	0	0	0	0	3
<i>Gymnetron veronicae</i> (Germar, 1821)	1	2	0	1	0	4
<i>Gymnetron villosulum</i> Gyllenhal, 1838	11	0	2	1	0	14
<i>Mecinus labilis</i> (Herbst, 1795)	1	5	1	1	0	8
<i>Mecinus pascuorum</i> (Gyllenhal, 1813)	31	25	71	7	2	136
<i>Mecinus pyraister</i> (Herbst, 1795)	24	8	41	7	1	81
<i>Miarus campanulae</i> (Linnaeus, 1767)	2	2	10	0	0	14
<i>Rhinusa antirrhini</i> (Paykull, 1800)	6	0	32	7	0	45
<i>Rhinusa linariae</i> (Panzer, 1795)	0	3	1	0	0	4
<i>Isochnus foliorum</i> (Müller, O.F., 1764)	0	2	2	1	1	6
<i>Orchestes (Orchestes) alni</i> (Linnaeus, 1758)	8	0	2	3	0	13
<i>Orchestes (Orchestes) pilosus</i> (Fabricius, 1781)	1	7	2	0	0	10
<i>Orchestes (Orchestes) quercus</i> (Linnaeus, 1758)	24	24	33	12	0	93
<i>Orchestes (Orchestes) rusci</i> (Herbst, 1795)	17	17	63	8	2	107
<i>Orchestes (Orchestes) signifer</i> (Creutzer, 1799)	8	7	11	5	0	31
<i>Orchestes (Orchestes) testaceus</i> (Müller, O.F., 1776)	0	0	1	0	0	1
<i>Orchestes (Salius) fagi</i> (Linnaeus, 1758)	14	31	109	34	2	190
<i>Pseudorchestes pratensis</i> (Germar, 1821)	0	0	1	1	0	2
<i>Rhamphus oxyacanthae</i> (Marsham, 1802)	6	4	31	3	1	45
<i>Rhamphus pulicarius</i> (Herbst, 1795)	10	16	44	3	2	75
<i>Tachyerges salicis</i> (Linnaeus, 1758)	4	15	12	6	1	38
<i>Tachyerges stigma</i> (Germar, 1821)	0	1	1	1	0	3
<i>Smicronyx reichi</i> (Gyllenhal, 1836)	0	0	0	1	0	1
<i>Orthochaetes setiger</i> (Beck, 1817)	4	5	3	2	0	14
<i>Sibinia primita</i> (Herbst, 1795)	0	0	3	0	0	3
<i>Tychius lineatulus</i> Stephens, 1831	1	1	0	0	0	2
<i>Tychius meliloti</i> Stephens, 1831	7	2	17	1	0	27
<i>Tychius picirostris</i> (Fabricius, 1787)	38	23	169	22	11	263
<i>Tychius schneideri</i> (Herbst, 1795)	1	1	0	0	0	2
<i>Bagous (Bagous) limosus</i> (Gyllenhal, 1827)	0	1	4	1	0	6
<i>Bagous (Bagous) lutulosus</i> (Gyllenhal, 1827)	1	0	0	0	0	1
<i>Bagous (Bagous) subcarinatus</i> Gyllenhal, 1836	0	1	0	0	0	1
<i>Bagous (Bagous) tempestivus</i> (Herbst, 1795)	2	1	1	1	0	5
<i>Bagous (Ephimeropus) petro</i> (Herbst, 1795)	0	0	0	3	0	3
<i>Bagous (Hydronomus) alismatis</i> (Marsham, 1802)	3	1	13	2	0	19
<i>Aulacobaris lepidii</i> (Germar, 1824)	0	3	0	1	0	4
<i>Aulacobaris picicornis</i> (Marsham, 1802)	0	0	1	0	0	1
<i>Melanobaris laticollis</i> (Marsham, 1802)	0	1	0	0	0	1
<i>Limnobaris dolorosa</i> (Goeze, 1777)	10	14	16	8	0	48
<i>Limnobaris t-album</i> (Linnaeus, 1758)	0	6	2	3	1	12
<i>Amalorrhynchus melanarius</i> (Stephens, 1831)	0	0	9	0	0	9
<i>Calosirus terminatus</i> (Herbst, 1795)	1	0	0	0	0	1
<i>Ceutorhynchus alliariae</i> Brisout, 1860	0	2	3	2	0	7
<i>Ceutorhynchus assimilis</i> (Paykull, 1792)	133	65	201	47	11	457

Yorkshire Coleoptera checklist 2011

<i>Ceutorhynchus atomus</i> Boheman, 1845	1	0	0	1	0	2
<i>Ceutorhynchus chalybaeus</i> Germar, 1824	0	0	1	1	0	2
<i>Ceutorhynchus cochleariae</i> (Gyllenhal, 1813)	3	2	2	2	0	9
<i>Ceutorhynchus constrictus</i> (Marsham, 1802)	1	4	3	1	0	9
<i>Ceutorhynchus contractus</i> (Marsham, 1802)	20	15	19	13	2	69
<i>Ceutorhynchus erysimi</i> (Fabricius, 1787)	22	26	32	28	5	113
<i>Ceutorhynchus hirtulus</i> Germar, 1824	0	1	0	1	0	2
<i>Ceutorhynchus insularis</i> Dieckmann, 1971	0	0	1	0	0	1
<i>Ceutorhynchus pallidactylus</i> (Marsham, 1802)	68	24	132	31	19	274
<i>Ceutorhynchus parvulus</i> Brisout, Ch., 1869	0	0	1	0	0	1
<i>Ceutorhynchus picitarsis</i> Gyllenhal, 1837	8	0	10	0	0	18
<i>Ceutorhynchus pumilio</i> (Gyllenhal, 1827)	1	0	0	0	0	1
<i>Ceutorhynchus pyrrhorhynchus</i> (Marsham, 1802)	5	2	9	1	0	17
<i>Ceutorhynchus resedae</i> (Marsham, 1802)	2	0	2	0	0	4
<i>Ceutorhynchus sulcicollis</i> (Paykull, 1800)	1	0	0	1	0	2
<i>Ceutorhynchus turbatus</i> Schultze, 1903	1	0	0	0	0	1
<i>Ceutorhynchus typhae</i> (Herbst, 1795)	59	41	47	24	11	182
<i>Coeliodes rana</i> (Fabricius, 1787)	6	18	20	6	0	50
<i>Coeliodes ruber</i> (Marsham, 1802)	0	0	1	0	0	1
<i>Coeliodes transversealbofasciatus</i> (Goeze, 1777)	1	1	0	0	0	2
<i>Coeliodinus rubicundus</i> (Herbst, 1795)	8	18	30	7	0	63
<i>Datonychus angulosus</i> (Boheman, 1845)	1	0	1	0	0	2
<i>Datonychus melanostictus</i> (Marsham, 1802)	0	0	5	3	0	8
<i>Datonychus urticae</i> (Boheman, 1845)	0	0	1	0	0	1
<i>Drupenatus nasturtii</i> (Germar, 1824)	6	0	10	0	0	16
<i>Glocianus distinctus</i> (Brisout, Ch., 1870)	0	0	2	0	0	2
<i>Glocianus punctiger</i> (Gyllenhal, 1837)	1	0	1	0	0	2
<i>Hadroplontus litura</i> (Fabricius, 1775)	16	10	44	11	5	86
<i>Hadroplontus trimaculatus</i> (Fabricius, 1775)	0	3	0	0	0	3
<i>Micrelus ericae</i> (Gyllenhal, 1813)	4	18	42	4	6	74
<i>Microplontus campestris</i> (Gyllenhal, 1837)	1	0	10	1	0	12
<i>Microplontus rugulosus</i> (Herbst, 1795)	17	1	12	3	0	33
<i>Mogulones asperifoliarum</i> (Gyllenhal, 1813)	12	5	7	2	2	28
<i>Nedys quadrimaculatus</i> (Linnaeus, 1758)	62	33	213	65	15	388
<i>Parethelcus pollinarius</i> (Forster, 1771)	36	19	91	15	0	161
<i>Poophagus sisymbrii</i> (Fabricius, 1777)	9	7	12	12	1	41
<i>Sirocalodes depressicollis</i> (Gyllenhal, 1813)	2	1	0	3	0	6
<i>Sirocalodes mixtus</i> (Mulsant & Rey, 1858)	2	0	1	0	0	3
<i>Sirocalodes quercicola</i> (Paykull, 1792)	0	0	1	0	0	1
<i>Stenocarus ruficornis</i> (Stephens, 1831)	2	7	0	2	0	11
<i>Thamiocolus viduatus</i> (Gyllenhal, 1813)	1	0	0	0	0	1
<i>Trichosirocalus dawsoni</i> (Brisout, Ch., 1869)	0	1	0	0	0	1
<i>Trichosirocalus horridus</i> (Panzer, 1801)	0	2	1	0	0	3
<i>Trichosirocalus thalhammeri</i> (Schultze, 1906)	1	0	0	0	0	1
<i>Trichosirocalus troglodytes</i> (Fabricius, 1787)	53	37	100	14	0	204
<i>Zacladus exiguus</i> (Olivier, 1807)	0	0	0	1	0	1
<i>Zacladus geranii</i> (Paykull, 1800)	13	9	14	20	11	67
<i>Amalus scortillum</i> (Herbst, 1795)	7	1	7	5	0	20
<i>Eubrychius velutus</i> (Beck, 1817)	0	0	6	2	0	8
<i>Neophytobius muricatus</i> (Brisout, Ch., 1867)	0	1	0	0	0	1

Yorkshire Coleoptera checklist 2011

<i>Neophytobius quadrinodosus</i> (Gyllenhal, 1813)	1	2	0	0	0	3
<i>Pelenomus canaliculatus</i> (Fåhraeus, 1843)	0	1	1	1	0	3
<i>Pelenomus comari</i> (Herbst, 1795)	2	3	2	8	0	15
<i>Pelenomus quadricorniger</i> (Colonnelli, 1986)	0	0	1	0	0	1
<i>Pelenomus quadrituberculatus</i> (Fabricius, 1787)	9	8	13	10	0	40
<i>Phytobius leucogaster</i> (Marsham, 1802)	1	2	3	0	0	6
<i>Rhinoncus bruchoides</i> (Herbst, 1784)	2	1	0	0	0	3
<i>Rhinoncus castor</i> (Fabricius, 1792)	16	8	27	6	2	59
<i>Rhinoncus inconspectus</i> (Herbst, 1795)	17	1	10	2	0	30
<i>Rhinoncus pericarpus</i> (Linnaeus, 1758)	49	26	108	29	9	221
<i>Rhinoncus perpendicularis</i> (Reich, 1797)	25	6	31	10	2	74
<i>Rhopalomesites tardyi</i> (Curtis, 1825)	0	7	3	1	0	11
<i>Pselactus spadix</i> (Herbst, 1795)	17	0	0	0	0	17
<i>Pseudophloeophagus aeneopiceus</i> (Boheman, 1845)	0	1	0	0	0	1
<i>Euophryum confine</i> (Broun, 1881)	17	7	59	7	0	90
<i>Euophryum rufum</i> (Broun, 1880)	0	0	1	1	0	2
<i>Pentarthrum huttoni</i> Wollaston, 1854	2	0	8	2	0	12
<i>Rhyncolus ater</i> (Linnaeus, 1758)	0	2	2	0	0	4
<i>Phloeophagus lignarius</i> (Marsham, 1802)	2	0	8	1	0	11
<i>Cryptorhynchus lapathi</i> (Linnaeus, 1758)	1	3	18	3	2	27
<i>Acalles misellus</i> Boheman, 1844	5	2	3	0	0	10
<i>Acalles ptinoides</i> (Marsham, 1802)	0	2	14	2	4	22
<i>Kykliaacalles roboris</i> (Curtis, 1835)	0	7	9	4	0	20
<i>Gronops lunatus</i> (Fabricius, 1775)	1	0	4	0	0	5
<i>Graptus triguttatus</i> (Fabricius, 1775)	10	7	1	9	3	30
<i>Neliocarus faber</i> (Herbst, 1784)	1	3	3	1	0	8
<i>Neliocarus nebulosus</i> (Stephens, 1831)	2	20	12	4	0	38
<i>Neliocarus sus</i> (Stephens, 1831)	11	27	16	4	3	61
<i>Strophosoma capitatum</i> (De Geer, 1775)	1	2	5	3	0	11
<i>Strophosoma melanogrammum</i> (Forster, 1771)	26	42	146	17	6	237
<i>Attactagenus plumbeus</i> (Marsham, 1802)	6	9	7	1	1	24
<i>Philopeton plagiatum</i> (Schaller, 1783)	37	5	5	0	0	47
<i>Barynotus moerens</i> (Fabricius, 1792)	2	12	27	11	4	56
<i>Barynotus obscurus</i> (Fabricius, 1775)	8	11	30	8	2	59
<i>Barynotus squamosus</i> Germar, 1824	0	4	10	8	5	27
<i>Omiamima mollina</i> (Boheman, 1834)	10	5	15	2	0	32
<i>Otiorhynchus (Otiorhynchus) clavipes</i> (Bonsdorff, 1785)	0	1	0	0	0	1
<i>Otiorhynchus (Otiorhynchus) ligneus</i> (Olivier, 1807)	1	1	2	3	0	7
<i>Otiorhynchus (Otiorhynchus) nodosus</i> (Müller, O. F. 1764)	0	4	3	3	2	12
<i>Otiorhynchus (Otiorhynchus) porcatus</i> (Herbst, 1795)	0	2	2	3	0	7
<i>Otiorhynchus (Otiorhynchus) raucus</i> (Fabricius, 1777)	0	1	1	2	0	4
<i>Otiorhynchus (Otiorhynchus) rugosostriatus</i> (Goeze, 1777)	2	1	10	1	0	14
<i>Otiorhynchus (Arammichnus) atroapterus</i> (De Geer, 1775)	0	3	1	0	0	4
<i>Otiorhynchus (Arammichnus) ligustici</i> (Linnaeus, 1758)	0	0	1	0	0	1
<i>Otiorhynchus (Dorymerus) rugifrons</i> (Gyllenhal, 1813)	1	3	4	1	3	12
<i>Otiorhynchus (Dorymerus) singularis</i> (Linnaeus, 1767)	31	52	198	31	12	324
<i>Otiorhynchus (Dorymerus) sulcatus</i> (Fabricius, 1775)	5	13	70	11	6	105
<i>Otiorhynchus (Tournieria) desertus</i> Rosenhauer, 1847	0	0	2	0	1	3
<i>Otiorhynchus (Tournieria) ovatus</i> (Linnaeus, 1758)	39	6	19	9	1	74
<i>Caenopsis fissirostris</i> (Walton, 1847)	0	0	0	1	0	1

Yorkshire Coleoptera checklist 2011

Phyllobius (Phyllobius) pyri (Linnaeus, 1758)	57	59	267	38	19	440
Phyllobius (Phyllobius) vespertinus (Fabricius, 1792)	0	3	1	0	0	4
Phyllobius (Dieletus) argentatus (Linnaeus, 1758)	16	44	192	24	17	293
Phyllobius (Metaphyllobius) glaucus (Scopoli, 1763)	6	41	109	18	9	183
Phyllobius (Metaphyllobius) pomaceus Gyllenhal, 1834	17	30	117	28	5	197
Phyllobius (Nemoicus) oblongus (Linnaeus, 1758)	7	20	135	27	12	201
Phyllobius (Parnemoicus) roboretanus Gredler, 1882	63	49	202	39	13	366
Phyllobius (Parnemoicus) viridicollis (Fabricius, 1801)	5	7	2	21	1	36
Phyllobius (Pterygorrhynchus) maculicornis Germar, 1824	8	16	76	13	0	113
Phyllobius (Subphyllobius) virideaeris (Laicharting, 1781)	2	0	3	0	0	5
Liophloeus tessulatus (Müller, O.F., 1776)	22	11	32	7	2	74
Polydrusus (Polydrusus) tereticollis (De Geer, 1775)	0	29	58	5	0	92
Polydrusus (Chrysophis) formosus (Mayer, 1779)	0	0	2	0	0	2
Polydrusus (Eudipnus) mollis (Ström, 1768)	0	27	9	4	4	44
Polydrusus (Eurodrusus) confluens Stephens, 1831	1	3	2	0	0	6
Polydrusus (Eustolus) flavipes (De Geer, 1775)	1	0	0	0	0	1
Polydrusus (Eustolus) pterygomalis Boheman, 1840	4	43	19	15	4	85
Polydrusus (Neoeustolus) cervinus (Linnaeus, 1758)	15	27	80	10	4	136
Polydrusus (Neoeustolus) pilosus Gredler, 1866	1	9	3	2	2	17
Polydrusus (Neoeustolus) pulchellus Stephens, 1831	3	0	0	0	0	3
Barypeithes (Barypeithes) sulcifrons (Boheman, 1843)	2	3	7	0	0	12
Barypeithes (Exomias) araneiformis (Schrank, 1781)	8	22	41	18	5	94
Barypeithes (Exomias) pellucidus (Boheman, 1834)	26	43	205	41	11	326
Barypeithes (Exomias) pyrenaeus Seidlitz, 1868	0	0	3	0	0	3
Brachysomus echinatus (Bonsdorff, 1785)	5	11	8	10	1	35
Sciaphilus asperatus (Bonsdorff, 1785)	4	10	42	15	2	73
Sitona (Sitona) ambiguus Gyllenhal, 1834	0	3	1	1	0	5
Sitona (Sitona) cambricus Stephens, 1831	0	0	7	1	0	8
Sitona (Sitona) cylindricollis (Fåhraeus, 1840)	8	2	7	0	1	18
Sitona (Sitona) hispidulus (Fabricius, 1777)	21	22	66	19	8	136
Sitona (Sitona) humeralis Stephens, 1831	10	3	16	8	1	38
Sitona (Sitona) lepidus Gyllenhal, 1834	24	34	93	30	12	193
Sitona (Sitona) lineatus (Linnaeus, 1758)	163	95	395	101	26	780
Sitona (Sitona) lineellus (Bonsdorff, 1785)	5	3	3	2	0	13
Sitona (Sitona) macularius (Marsham, 1802)	2	2	0	0	0	4
Sitona (Sitona) ononidis Sharp, 1866	0	1	2	0	0	3
Sitona (Sitona) puncticollis Stephens, 1831	9	10	8	3	0	30
Sitona (Sitona) regensteinensis (Herbst, 1797)	13	9	89	7	3	121
Sitona (Sitona) striatellus Gyllenhal, 1834	10	3	13	6	1	33
Sitona (Sitona) sulcifrons (Thunberg, 1798)	6	7	9	9	2	33
Sitona (Sitona) suturalis Stephens, 1831	20	23	54	15	5	117
Sitona (Sitona) waterhousei Walton, 1846	0	1	1	0	0	2
Sitona (Charagmus) griseus (Fabricius, 1775)	3	2	5	0	0	10
Tanymecus palliatus (Fabricius, 1787)	2	5	1	1	0	9
Trachyploeus alternans Gyllenhal, 1834	0	2	0	0	0	2
Trachyploeus angustisetulus Hansen, 1915	0	0	1	0	0	1
Trachyploeus aristatus (Gyllenhal, 1827)	1	1	4	0	0	6
Trachyploeus asperatus Boheman, 1843	0	2	1	0	0	3
Trachyploeus bifoveolatus (Beck, 1817)	1	2	3	0	0	6
Trachyploeus scabriculus (Linnaeus, 1771)	2	0	1	0	0	3

Yorkshire Coleoptera checklist 2011

<i>Cathormiocerus myrmecophilus</i> (Seidlitz, 1868)	1	0	0	0	0	1
<i>Tropiphorus elevatus</i> (Herbst, 1795)	0	1	2	1	0	4
<i>Tropiphorus obtusus</i> (Bonsdorff, 1785)	0	1	1	3	0	5
<i>Tropiphorus terricola</i> (Newman, 1838)	2	7	8	2	5	24
<i>Hypera</i> (<i>Hypera</i>) <i>arator</i> (Linnaeus, 1758)	10	2	5	3	0	20
<i>Hypera</i> (<i>Hypera</i>) <i>fuscocinerea</i> (Marsham, 1802)	0	2	0	0	0	2
<i>Hypera</i> (<i>Hypera</i>) <i>nigrirostris</i> (Fabricius, 1775)	27	10	38	15	0	90
<i>Hypera</i> (<i>Hypera</i>) <i>plantaginis</i> (De Geer, 1775)	2	8	11	9	0	30
<i>Hypera</i> (<i>Hypera</i>) <i>postica</i> (Gyllenhal, 1813)	17	9	12	6	0	44
<i>Hypera</i> (<i>Hypera</i>) <i>suspiciosa</i> (Herbst, 1795)	1	3	5	3	3	15
<i>Hypera</i> (<i>Hypera</i>) <i>venusta</i> (Fabricius, 1781)	2	1	5	0	0	8
<i>Hypera</i> (<i>Antidonus</i>) <i>dauci</i> (Olivier, 1807)	3	1	0	0	0	4
<i>Hypera</i> (<i>Antidonus</i>) <i>zoilus</i> (Scopoli, 1763)	7	8	5	11	0	31
<i>Hypera</i> (<i>Boreohypera</i>) <i>diversipunctata</i> (Schrank, 1798)	3	0	1	2	0	6
<i>Hypera</i> (<i>Dapalinus</i>) <i>meles</i> (Fabricius, 1792)	0	0	1	1	0	2
<i>Hypera</i> (<i>Eirinomorphus</i>) <i>pollux</i> (Fabricius, 1801)	10	2	3	3	0	18
<i>Hypera</i> (<i>Eirinomorphus</i>) <i>rumicis</i> (Linnaeus, 1758)	4	3	7	4	1	19
<i>Limobius borealis</i> (Paykull, 1792)	2	3	0	5	2	12
<i>Lixus</i> (<i>Lixus</i>) <i>paraplecticus</i> (Linnaeus, 1758)	1	0	1	0	0	2
<i>Lixus</i> (<i>Dilixellus</i>) <i>angustatus</i> (Fabricius, 1775)	0	1	0	0	0	1
<i>Cleonis pigra</i> (Scopoli, 1763)	15	2	0	0	0	17
<i>Magdalis</i> (<i>Magdalis</i>) <i>duplicata</i> Germar, 1818	0	0	0	1	0	1
<i>Magdalis</i> (<i>Magdalis</i>) <i>phlegmatica</i> (Herbst, 1797)	0	0	0	2	0	2
<i>Magdalis</i> (<i>Edo</i>) <i>ruficornis</i> (Linnaeus, 1758)	3	5	6	6	1	21
<i>Magdalis</i> (<i>Odontomagdalis</i>) <i>armigera</i> (Fourcroy, 1785)	3	7	12	7	2	31
<i>Magdalis</i> (<i>Odontomagdalis</i>) <i>carbonaria</i> (Linnaeus, 1758)	2	3	6	4	1	16
<i>Magdalis</i> (<i>Panus</i>) <i>barbicornis</i> (Latreille, 1804)	1	0	3	0	0	4
<i>Magdalis</i> (<i>Porrothus</i>) <i>cerasi</i> (Linnaeus, 1758)	3	1	11	5	0	20
<i>Liparus coronatus</i> (Goeze, 1777)	0	1	0	0	0	1
<i>Leiosoma deflexum</i> (Panzer, 1795)	11	25	132	21	12	201
<i>Leiosoma oblongulum</i> Boheman, 1842	0	0	2	2	0	4
<i>Anoplus plantaris</i> (Naezen, 1794)	4	16	26	2	0	48
<i>Anoplus roboris</i> Suffrian, 1840	0	1	2	1	0	4
<i>Hylobius</i> (<i>Callirus</i>) <i>abietis</i> (Linnaeus, 1758)	6	5	11	6	1	29
<i>Pissodes castaneus</i> (De Geer, 1775)	3	2	1	1	0	7
<i>Pissodes pini</i> (Linnaeus, 1758)	4	6	9	3	0	22
<i>Trachodes hispidus</i> (Linnaeus, 1758)	0	0	1	0	0	1
<i>Orobitis cyaneus</i> (Linnaeus, 1758)	1	7	1	2	0	11
<i>Scolytus intricatus</i> (Ratzeburg, 1837)	1	9	10	7	0	27
<i>Scolytus mali</i> (Bechstein & Scharfenberg, 1805)	0	3	3	4	0	10
<i>Scolytus multistriatus</i> (Marsham, 1802)	1	0	17	0	1	19
<i>Scolytus ratzeburgi</i> Janson, 1856	0	1	1	0	0	2
<i>Scolytus rugulosus</i> (Müller, P.W.J., 1818)	1	0	3	2	0	6
<i>Scolytus scolytus</i> (Fabricius, 1775)	1	9	32	7	0	49
<i>Pityophthorus lichtensteinii</i> (Ratzeburg, 1837)	1	0	2	0	0	3
<i>Pityophthorus pubescens</i> (Marsham, 1802)	1	1	0	0	0	2
<i>Cryphalus asperatus</i> (Gyllenhal, 1813)	2	5	0	3	0	10
<i>Ernoporicus caucasicus</i> (Lindemann, 1876)	0	2	0	1	0	3
<i>Ernoporicus fagi</i> (Fabricius, 1798)	0	0	0	2	0	2
<i>Ernoporus tiliae</i> (Panzer, 1793)	0	0	1	1	0	2

Yorkshire Coleoptera checklist 2011

<i>Dryocoetes alni</i> (Georg, 1856)	0	0	5	5	2	12
<i>Dryocoetes autographus</i> (Ratzeburg, 1837)	2	7	0	4	0	13
<i>Dryocoetes villosus</i> (Fabricius, 1792)	1	8	23	6	0	38
<i>Taphrorhynchus bicolor</i> (Herbst, 1793)	0	0	0	1	0	1
<i>Ips acuminatus</i> (Gyllenhal, 1827)	1	1	0	0	0	2
<i>Ips sexdentatus</i> (Boerner, 1776)	0	0	1	0	0	1
<i>Ips typographus</i> (Linnaeus, 1758)	1	1	0	0	0	2
<i>Orthotomicus laricis</i> (Fabricius, 1792)	2	1	0	1	0	4
<i>Pityogenes bidentatus</i> (Herbst, 1783)	5	7	9	3	1	25
<i>Pityogenes chalcographus</i> (Linnaeus, 1761)	1	4	1	0	0	6
<i>Pityogenes quadridens</i> (Hartig, 1834)	0	0	2	0	0	2
<i>Xyleborinus saxesenii</i> (Ratzeburg, 1837)	2	4	4	1	0	11
<i>Xyleborus dispar</i> (Fabricius, 1792)	0	1	6	1	0	8
<i>Xyleborus dryographus</i> (Ratzeburg, 1837)	0	0	2	0	0	2
<i>Trypodendron domesticum</i> (Linnaeus, 1758)	2	10	26	11	1	50
<i>Trypodendron lineatum</i> (Olivier, 1795)	0	1	4	1	0	6
<i>Trypodendron signatum</i> (Fabricius, 1792)	0	3	6	0	0	9
<i>Hylesinus crenatus</i> (Fabricius, 1787)	1	6	12	5	3	27
<i>Hylesinus toranio</i> (Danthoine in Bernard, 1788)	3	2	5	2	0	12
<i>Hylesinus varius</i> (Fabricius, 1775)	5	9	25	6	4	49
<i>Hylastinus obscurus</i> (Marsham, 1802)	1	7	7	0	1	16
<i>Kissophagus hederæ</i> (Schmitt, 1843)	0	2	0	0	0	2
<i>Pteleobius vittatus</i> (Fabricius, 1787)	1	0	4	1	0	6
<i>Hylastes ater</i> (Fabricius, 1792)	2	5	7	2	1	17
<i>Hylastes brunneus</i> Erichson, 1836	0	1	1	0	1	3
<i>Hylastes cunicularius</i> Erichson, 1836	0	2	0	1	0	3
<i>Hylastes opacus</i> Erichson, 1836	2	2	2	0	0	6
<i>Hylurgops palliatus</i> (Gyllenhal, 1813)	7	9	12	3	0	31
<i>Phloeotribus rhododactylus</i> (Marsham, 1802)	7	4	6	0	0	17
<i>Polygraphus poligraphus</i> (Linnaeus, 1758)	0	1	0	0	0	1
<i>Tomicus piniperda</i> (Linnaeus, 1758)	2	8	10	3	0	23
<i>Xylechinus pilosus</i> (Ratzeburg, 1837)	0	1	0	0	0	1
<i>Tanysphyrus lemnae</i> (Paykull, 1792)	12	7	24	13	0	56

Platypodidae [1/1]

<i>Platypus cylindrus</i> (Fabricius, 1792)	1	1	2	0	0	4
---	---	---	---	---	---	---

It has not been possible to include a column containing species status in this version of the checklist owing to software problems.

Please pass comments on the content of this checklist either to R J Marsh or M L Denton.